



Food and Agriculture Organization  
of the United Nations

ISSN 2709-006X [Print]  
ISSN 2709-0078 [Online]



FAOSTAT ANALYTICAL BRIEF 50

# Greenhouse gas emissions from agrifood systems

Global, regional and country trends, 2000–2020

## HIGHLIGHTS

- In 2020, global agrifood systems emissions were 16 billion tonnes of carbon dioxide equivalent (Gt CO<sub>2</sub>eq), an increase of 9 percent since 2000.
- Per capita emissions decreased by 15 percent over the same period, to 2.0 t CO<sub>2</sub>eq per capita in 2020.
- The share of agrifood systems emissions in emissions from all sectors dropped from 38 percent in 2000 to 31 percent in 2020, due to much faster growth in non-food emissions.
- Globally, the farm gate in 2020 represented nearly half of total agrifood systems emissions, pre- and post-production processes contributed one-third and land-use change one-fifth.
- In 2020, at the regional level, farm-gate emissions were the largest component in Oceania (71 percent), Asia (50 percent) and the Americas (43 percent). Land-use change was the largest contributor in Africa (44 percent), while pre- and post-production processes were the largest contributor in Europe (53 percent).
- Per capita emissions were the highest in Oceania (6.5 t CO<sub>2</sub>eq per capita) and the lowest in Asia (1.4 t CO<sub>2</sub>eq per capita). Agrifood systems accounted for the largest share in emissions from all sectors in Africa (59 percent) and the lowest in Asia (near 25 percent).
- Globally, emissions intensities in 2020 varied between 1 kg CO<sub>2</sub>eq/kg and 30 kg CO<sub>2</sub>eq/kg for meat (with the lowest values for chicken meat and the highest for beef). The global farm-gate emissions intensity for cow milk was 1 kg CO<sub>2</sub>eq/kg, about 24 percent less than in 2000.

## FAOSTAT EMISSIONS

### BACKGROUND

Agrifood systems account for one-third of total anthropogenic greenhouse gas (GHG) emissions (Crippa *et al.*, 2021; Tubiello *et al.*, 2021). They are generated within the *farm gate*, by crop and livestock production activities; by *land-use change*, for instance deforestation and peatland drainage to make room for agriculture; and in *pre- and post-production processes*, such as food manufacturing, retail, household consumption and food disposal (Tubiello *et al.*, 2022).

Statistics on underlying activity data, emissions and indicators (emissions per capita, shares in total emissions, and emissions per commodity) are disseminated in FAOSTAT at country, regional and global levels, covering over 200 countries and territories, for the period 1961–2020 (FAO, 2022a, 2022b,



2022c), and expressed in both single component gases – carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and fluorinated gases (F-gases) – and their cumulative carbon dioxide equivalents (CO<sub>2</sub>eq). This analysis focuses on results relative to the period 2000–2020.


## GLOBAL

In 2020, global annual anthropogenic GHG emissions reached 52 Gt CO<sub>2</sub>eq, down 4 percent from 54 Gt CO<sub>2</sub>eq in 2019 – reflecting a well-documented reduction in economic activities due to the COVID-19 pandemic. They were nonetheless 34 percent higher than in 2000. At the same time, emissions from agrifood systems were 16 Gt CO<sub>2</sub>eq in 2020, down 3 percent from 2019, but 9 percent higher than in 2000. The share of agrifood systems in total emissions in 2020 (31 percent) confirmed the steady downward trend from the levels of 2000 (38 percent), a consequence of agrifood systems emissions growing significantly more slowly than the rest of the economy, dominated by fossil fuels combustion for energy use. In fact, non-food emissions grew nearly 50 percent since 2000. Agrifood systems emissions per capita likewise decreased over the period, from 2.4 t CO<sub>2</sub>eq/cap to 2.0 t CO<sub>2</sub>eq/cap (Figure 1).

Among the three components of agrifood systems in 2020, farm-gate emissions were nearly half of the total (7.4 Gt CO<sub>2</sub>eq), followed by emissions from pre- and post-production (5.6 Gt CO<sub>2</sub>eq) and land-use change (3.1 Gt CO<sub>2</sub>eq). Over the period 2000-2020, pre- and post-production emissions grew the fastest (45 percent); emissions within the farm gate increased by 13 percent while those from land-use change significantly decreased (-29 percent), largely a result of the long-term slowdown in deforestation rates (Figure 1). With respect to 2019, land-use change emissions in 2020 decreased significantly (-11 percent) due to lower fire intensity in degraded tropical peatlands. Pre- and post-production emissions also decreased (-4 percent), in line with the COVID-19 decrease in fossil fuel energy use. Conversely, emissions from the farm gate increased 1 percent in 2020, consistently with recent FAOSTAT statistics of crop and livestock production, which had indicated the limited impact of the COVID-19 pandemic at the global level on this primary sector.



Figure 1: Global agrifood systems emissions by component and indicator




**Source:** FAO. 2022. Emissions totals. In: FAO. Rome. Cited October 2022. <https://www.fao.org/faostat/en/#data/GT> and FAO. 2022. Emissions shares. In: FAO. Rome. Cited October 2022. <https://www.fao.org/faostat/en/#data/EM>

When breaking down agrifood systems emissions by gas, CO<sub>2</sub> emissions remained stable to 7.9 Gt as the reduced emissions from deforestation were cancelled out by increased CO<sub>2</sub> emissions in pre- and post-production processes; CH<sub>4</sub> emissions grew from 173 Mt to 193 Mt (+14 percent) and N<sub>2</sub>O emissions increased from 7 Mt to 9 Mt (+12 percent). Over the past two decades, emissions from agrifood systems decreased from 29 percent to 21 percent of the total for CO<sub>2</sub>, from 58 percent to 53 percent for CH<sub>4</sub>, while the share of N<sub>2</sub>O agrifood systems emissions remained at 78 percent over the entire period. Finally, the share of F-gases generated by cold chains, largely in food retail, decreased from 32 percent to 26 percent.

A further breakdown by subcomponent helps to highlight the relative importance of specific processes across global production, supply and consumption chains. In 2020, the most important contributors to global agrifood systems emissions were CO<sub>2</sub> emissions from deforestation (2.9 Gt CO<sub>2</sub>eq) and CH<sub>4</sub> from enteric fermentation of ruminant livestock (2.8 Gt CO<sub>2</sub>eq), representing together nearly 40 percent of the total. Other important global contributors were CH<sub>4</sub> emissions from livestock manure and food systems waste disposal, and CO<sub>2</sub> emissions from household consumption, at about 1.3 Gt CO<sub>2</sub>eq each. All these components are shown in Figure 2.

**Figure 2: Detailed composition of agrifood systems emissions (2020)**




**Note:** Emissions/removals on forestland (which are not part of agrifood systems emissions) are also shown as included in the FAOSTAT Emissions database.

**Source:** FAO. 2022. Emissions totals. In: FAO. Rome. Cited October 2022. <https://www.fao.org/faostat/en/#data/GT>

Emissions intensities, defined as the GHG emissions within the farm gate per unit weight of product (kg CO<sub>2</sub>eq/kg product) were computed for several primary commodities: six types of meat (buffalo, cattle [beef], chicken, goat, pig and sheep), four types of milk (camel, cow, goat and sheep), hen eggs, rice, and other cereals (combining the intensities of barley, maize, millet, oats, rye, sorghum and wheat). Figure 3 focuses on a subset of them. In 2020, farm-gate emissions by kg of beef were 32 kg CO<sub>2</sub>eq/kg, a high value that is largely due to methane production by ruminant fermentation. Indeed, the emissions intensities of monogastric animals were much smaller: nearly 2 kg CO<sub>2</sub>eq/kg for pork and less than 1 kg CO<sub>2</sub>eq/kg for chicken. The global emissions intensity of cow milk was 1 kg CO<sub>2</sub>eq per kg of milk. The global emissions intensities of cereals were 1 kg CO<sub>2</sub>eq/kg for rice and 0.2 kg CO<sub>2</sub>eq/kg for other cereals. The GHG intensity of hen eggs was 0.6 kg CO<sub>2</sub>eq/kg. Farm-gate emissions intensities had a marked long-term declining trend since 2000 across all commodities, with the largest reduction computed for cow milk (-24 percent) and rice (-14 percent) (Figure 3). Such reductions reflect increases in crop and livestock production efficiency over time, often achieved through economies of scale.



**Figure 3: Global emissions intensities (2020) and percentage change since 2000**



**Source:** FAO. 2022. Emissions intensities. In: FAO. Rome. Cited October 2022  
<https://www.fao.org/faostat/en/#data/EI>


## REGIONAL

In 2020, total agrifood systems emissions were the largest in Asia (6.6 Gt CO<sub>2</sub>eq) and the Americas (4.3 Gt CO<sub>2</sub>eq), mainly a reflection of area and population. The relative role of the three components in total emissions from agrifood systems varied across regions, reflecting structural differences in production and distribution systems around the world (Figure 4). Emissions in Africa and the Americas had significant land-use change components (1.2–1.3 Gt CO<sub>2</sub>eq), respectively 44 percent and 31 percent of the total agrifood systems emissions, reflecting the extensive nature of agriculture in both regions and its impact on surrounding ecosystems, mainly via deforestation. Conversely, significant pre- and post- production emissions were observed in Asia (43 percent, or 2.9 Gt CO<sub>2</sub>eq) and especially in Europe (53 percent or 1.0 Gt CO<sub>2</sub>eq), where this component was in fact the largest contributor. Emissions produced within the farm gate remained the dominant component of agrifood systems emissions in Oceania (71 percent), Asia (50 percent) and the Americas (42 percent). Furthermore, emissions from agrifood systems had significantly different trends across regions between 2000 and 2020. They decreased by 33 percent in Oceania, 9 percent in the Americas and 1 percent in Europe, while they grew by 35 percent in Africa and 20 percent in Asia.

Oceania was the largest per capita emitter (6.5 t CO<sub>2</sub>eq/cap) in 2020, followed by the Americas (4.2 t CO<sub>2</sub>eq/cap), respectively three and two times larger than the world average. Europe (2.7 t CO<sub>2</sub>eq/cap) and Africa (2.1 t CO<sub>2</sub>eq/cap) had values closer to the world average, while Asia was the lowest per capita emitter at 1.4 t CO<sub>2</sub>eq/cap.

Africa had the largest share of agrifood systems in total emissions (59 percent) in 2020, consistently with the predominance of agriculture in most economies of the region. The shares were much lower in the Americas (40 percent), Oceania (38 percent) and Europe (32 percent) and lowest in Asia (23 percent), reflecting the economic efficiency of more intensive mixed and modern production systems (Figure 4).

Figure 4: Regional agrifood systems emissions and share in total emissions (2020)



**Source:** FAO. 2022. Emissions totals. In: FAO. Rome. Cited October 2022. <https://www.fao.org/faostat/en/#data/GT> and FAO. 2022. Emissions shares. In: FAO. Rome. Cited October 2022. <https://www.fao.org/faostat/en/#data/EM>

The intensities of farm-gate emissions varied significantly among regions and followed different trends over the period depending on the commodity, as shown using the examples of beef and cow milk. These regional features generally reflected known differences across regions in farm production systems, from traditional to mixed systems to modern agriculture.

The 2020 emissions intensity of beef was highest in Africa (66 kg CO<sub>2</sub>eq/kg), followed by the Americas and Asia (29 ka CO<sub>2</sub>eq/ka in both regions). Oceania (21 ka CO<sub>2</sub>eq/ka) and Europe (17 ka CO<sub>2</sub>eq/ka)

预览已结束，完整报告链接和二维码如下：

[https://www.yunbaogao.cn/report/index/report?reportId=5\\_31872](https://www.yunbaogao.cn/report/index/report?reportId=5_31872)

