

# Carbon in Drylands: Desertification, Climate Change and Carbon Finance

A UNEP-UNDP-UNCCD Technical Note for Discussions at CRIC 7, Istanbul, Turkey - 03-14 November, 2008

Prepared on behalf of UNEP by UNEP-WCMC

Authors: Kate Trumper, Corinna Ravilious and Barney Dickson

31<sup>st</sup> October 2008

*Disclaimer: The contents of this report do not necessarily reflect the views or policies of UNEP-WCMC or contributory organisations. The designations employed and the presentations do not imply the expressions of any opinion whatsoever on the part of UNEP-WCMC or contributory organisations concerning the legal status of any country, territory, city or area or its authority, or concerning the delimitation of its frontiers or boundaries.*

# **Technical Note: Carbon in drylands - Desertification, climate change and carbon finance**

## **Introduction**

Drylands cover about 40% of the Earth's land surface, excluding Antarctica and Greenland, and are home to more than two billion people (WRI 2002). They are susceptible to desertification, land degradation and drought (DLDD) and their populations, agriculture and ecosystems are vulnerable to climate change and variability. The United Nations Convention to Combat Desertification (UNCCD), one of the three 'Rio' conventions born out of the 1992 United Nations Conference on Environment and Development (UNCED), aims to address these issues and emphasises action to promote sustainable development at the community level.


The other Rio conventions are the United Nations Framework Convention on Climate Change (UNFCCC) and the Convention on Biological Diversity (CBD). The areas of interest of the three Conventions are closely linked and each has accepted the need to work in concert. One area of joint interest is that of the uptake of carbon dioxide from the atmosphere by plants and its storage in ecosystems. It is perhaps the only practicable way of removing carbon dioxide from the atmosphere in the short term and therefore one of the few options for addressing its existing carbon load, as distinct to slowing future loading by reducing current and future emissions. Most attention so far has focussed on carbon sequestration by tropical forests. More recently, some have argued for a more holistic approach to terrestrial carbon (The Terrestrial Carbon Group, 2008). This paper reviews the potential for carbon sequestration in dryland ecosystems, which includes forests, but also covers other habitats, such as grasslands, and, importantly, soils. It also considers ways in which carbon storage in drylands affects land degradation issues.

## **Carbon storage in drylands**

Plants take up carbon dioxide from the atmosphere and incorporate it into plant biomass through photosynthesis. Some of this carbon is emitted back to the atmosphere but what is left—the live and the dead plant parts, above and below ground—make up an organic carbon reservoir. Some of the dead plant matter is incorporated into the soil in humus, thereby enhancing the soil organic carbon pool.

Plant biomass per unit area of drylands is low (about 6 kilograms per square meter) compared with many terrestrial ecosystems (about 10–18 kilograms). But the large surface area of drylands gives dryland carbon sequestration a global significance. In particular, total dryland soil organic carbon reserves comprise 27% of the global soil organic carbon reserves (MA 2005). The soil properties, such as the chemical composition of soil organic matter and the matrix in which it is held, determine the different capacities of the land to act as a store for carbon that has direct implications for capturing greenhouse gases (FAO 2004). The fact that many of the dryland soils have been degraded means that they are currently far from saturated with carbon and their potential to sequester carbon may be very high (Farage et al. 2003).

**Global carbon stock density in drylands (above and below ground biomass plus soil carbon)**



Carbon (tonnes/hectare)



Non dryland areas

- 1 North America  
2 Greenland  
3 Central America & Caribbean  
4 South America  
5 Europe  
6 North Eurasia  
7 Africa  
8 Middle East  
9 South Asia  
10 East Asia  
11 South East Asia  
12 Australia & New Zealand  
13 Pacific  
14 Antarctic & peripheral Islands

**Data sources:**

Ruesch, A. S. & H. K. Gibbs. Global biomass carbon stock map based on IPCC Tier-1 Methodology. Oak Ridge National Laboratory's Carbon Dioxide Information Analysis Center (in review).

GBP-DIS 2000, Global Soil Data Products CD-ROM, Global Soil Data Task, International Geosphere-Biosphere Programme, Data and Information System, Potsdam, Germany. Sourced from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. <http://www.daac.ornl.gov>

UNEP-WCMC, 2007. A spatial analysis approach to the global delineation of dryland areas of relevance to the CBD Programme of Work on Dry and Subhumid lands. [http://www.unep-wcmc.org/abslats/drylands/dryland\\_report\\_final\\_HR.pdf](http://www.unep-wcmc.org/abslats/drylands/dryland_report_final_HR.pdf)

The contents of the map presented here do not necessarily reflect the views or policies of UNEP-WCMC or contributory organizations. The designations employed and the presentations do not imply the expression of any opinion whatsoever on the part of UNEP-WCMC or contributory organizations concerning the legal status of any country, territory or area or its authority, or concerning the delimitation of its frontiers or boundaries.



© UNEP-VACMG, 2008

The map above shows how the density of carbon stored, that is, the mass of carbon per hectare, varies throughout drylands. The carbon densities are derived from two global datasets: the carbon stock in biomass is from a map based on IPCC Tier-1 Methodology using global land cover data. (Ruesch & Gibbs, in review); soil carbon is from Global Soil Data Products CD-ROM. (IGBP-DIS 2000). The delineation of drylands is from UNEP-WCMC's map of areas of relevance to the CBD's programme of work on dry and sub-humid lands (UNEP-WCMC 2007). The UNCCD defines drylands according to an aridity index: the ratio of mean annual precipitation to mean annual potential evapotranspiration. The CBD definition of 'drylands' used within its Programme of Work on Dry and Subhumid Lands (UNEP/CBD/SBSTTA/5/9) differs from the UNCCD definition described above in two ways:

- i. It includes hyperarid zones (CCD does not) (UNEP/CBD/SBSTTA/5/9), which represent approximately 6.6 percent of the Earth's land surface.
- ii. Major vegetation types are used to define dryland areas in addition to those defined according to the aridity index (UNEP/CBD/SBSTTA/5/9).

Table 1 gives a breakdown of the carbon stored in each region in drylands. Figures for the total carbon stock in each region are from Campbell *et al.* (2008) and are derived from the same data as the dryland figures. Estimates of carbon stored in each region are sensitive to changes in land cover type. Therefore for detailed regional or national purposes, it will be necessary to refine global land cover data with more detailed local data. Nevertheless, this global overview shows that dryland carbon storage accounts for more than one third of the global stock. In some regions, such as the Middle East and Africa, a very high proportion of carbon is in drylands, so any sequestration measures there would need to address dryland ecosystems. Even in regions such as Africa and South Asia, where moist forests contain a lot of carbon, dryland carbon storage is still significant.

**Table 1. Comparison of total and drylands carbon stocks in regions of the world**

Map number	Region	Total carbon stock per region (Gt)	Carbon stock in drylands(Gt)	Share of regional carbon stock held in drylands (%)
1	North America	388	121	31
2	Greenland	5	0	0
3	Central America & Caribbean	16	1	7
4	South America	341	115	34
5	Europe	100	18	18
6	North Eurasia	404	96	24
7	Africa	356	211	59
8	Middle East	44	41	94
9	South Asia	54	26	49
10	East Asia	124	41	33
11	South East Asia	132	3	2
12	Australia/New Zealand	85	68	80
13	Pacific	3	0	0
<b>Total</b>		<b>2053</b>	<b>743</b>	<b>36</b>

## Land degradation and carbon emissions

According to the Millennium Ecosystem Assessment, “some 10–20% of the world’s drylands suffer from one or more forms of land degradation. Despite the global concern aroused by desertification, the available data on the extent of land degradation in drylands (also called desertification) are extremely limited. In the early 1990s, the Global Assessment of Soil Degradation, based on expert opinion, estimated that 20% of drylands (excluding hyper-arid areas) were affected by soil degradation. A study based on regional data sets (including hyper-arid drylands) derived from literature reviews, erosion models, field assessments and remote sensing found much lower levels of land degradation in drylands. Coverage was not complete, but the main areas of degradation were estimated to cover 10% of global drylands.” The MA estimated that the true level of degradation lay somewhere between the 10% and 20% figures. (MA 2005). The Land Degradation Assessment in Drylands (LADA) project, funded by the Global Environmental Facility (GEF) and carried out by the Food and Agriculture Organization of the United Nations (FAO) is drawing together information about degradation and developing ways of assessing the extent of land degradation and its impacts.

Land use change and degradation are important sources of greenhouse gases globally, responsible for about 20% of emissions (IPCC, 2007). Land degradation leads to increased carbon emissions both through loss of biomass when vegetation is destroyed and through increased soil erosion. Erosion leads to emissions in two ways: by reducing primary productivity, thereby reducing soils’ potential to store carbon and through direct losses of stored organic matter. Although not all carbon in eroded soil is returned to the atmosphere immediately, the net effect of erosion is likely to be increased carbon emissions (MA, 2005).

There have been a number of estimates of the rate of carbon emissions due to land degradation in drylands at different scales. At the global scale, Lal (2001) estimated that dryland ecosystems contribute 0.23 – 0.29 Gt of carbon a year to the atmosphere, which is about 4% of global emissions from all sources combined (MA 2005). In China, degradation of grassland, particularly on the Qinghai-Tibetan Plateau, has led to the loss of 3.56 Gt soil organic carbon over the last 20 years. It is estimated that the soils of China overall now act as a net carbon source, with a loss of 2.86 Gt in the same period (Xie et al., 2007). It is therefore vital from a climate perspective that this region is managed to enhance carbon sequestration (Xu et al., 2004) and further study is clearly required in this area (ESPA China 2008).

Grace et al. (2006) reviewed carbon fluxes in tropical savannas. They found that carbon sequestration rates in these ecosystems may average 0.14 tonnes carbon per hectare per year or 0.39 tonnes carbon per hectare per year. They concluded that “if savannas were to be protected from fire and grazing, most of them would accumulate substantial carbon and the sink would be larger. Savannas are under anthropogenic pressure, but this has been much less publicized than deforestation in the rain forest biome. The rate of loss is not well established, but may exceed 1% per year, approximately twice as fast as that of rain forests. Globally, this is likely to constitute a flux to the atmosphere that is at least as large as that arising from deforestation of the rain forest.”

As well as contributing to greenhouse gas emissions, drylands are themselves vulnerable to the effects of climate change and the impacts of climate change in these areas may lead in turn to further carbon emissions. Any further failure of plant growth due to increased temperatures would further reduce carbon inputs to the soil, accelerating its degradation. Smith et al point out that ‘even partial loss of vegetation integrity could make soils more vulnerable to degradation through other agents such as grazing and cultivation.’ (Smith et al 2008)

## **Climate change mitigation through addressing DLDD**

Addressing land degradation in dryland ecosystems presents two complementary ways of mitigating climate change. First, by slowing or halting degradation, associated emissions can be similarly reduced. Second, and arguably of greater significance, changes in land management practices can lead to greater carbon sequestration, that is, to removing carbon from the atmosphere. In general, the carbon storage potential of dryland ecosystems is lower than for moist tropical systems, but the large area of drylands means that overall they have significant scope for sequestration.

### ***Managing drylands for carbon sequestration***

Since carbon losses from drylands are associated with loss of vegetation cover and soil erosion, management interventions that slow or reverse these processes can simultaneously achieve carbon sequestration. There is a wide range of strategies to increase the stock of carbon in the soil. Examples include enhancing soil quality, erosion control, afforestation and woodland regeneration, no-till farming, cover crops, nutrient management, manuring and sludge application, optimal livestock densities, water conservation and harvesting, efficient irrigation, land-use change (crops to grass/trees), set-aside, agroforestry, and the use of legumes (FAO 2004, Lal 2004, Smith 2008).

There is a growing interest in assessing the carbon sequestration potential of such strategies quantitatively. Using a modelling approach, Farage et al (2007) found the most effective practices for increasing soil carbon storage were those that maximised the input of organic matter, particularly farmyard manure (up to 0.09 tonnes C per hectare per year), maintaining trees (up to 0.15 tonnes C per hectare per year) and adopting zero tillage (up to 0.04 tonnes C per hectare per year (Farage et al 2007).

Tiessen et al. (1998) reviewed data on carbon and biomass budgets under different land use in tropical savannas and some dry forests in West Africa and North-Eastern Brazil. They found that improvements in the carbon sequestration in these semi arid regions depended on an increase in crop production under suitable rotations, improved fallow and animal husbandry, and a limitation on biomass burning. Use of fertilizer was required for improved productivities but socioeconomic constraints largely prevented such improvements, resulting in a very limited scope for changes in soil carbon management.

Increasing carbon stocks in the soil increases soil fertility, workability, water holding capacity, and reduces erosion risk and can thus reduce the vulnerability of managed soils to future global warming (Smith, 2008). However, hidden costs also need to be

considered, such as the addition of mineral or organic fertilizer (especially nitrogen and phosphorus) and water, which would need significant capital investment (MA 2005).

### ***Estimates of dryland carbon sequestration potential***

Several studies have attempted to assess the potential for carbon sequestration in drylands. Considering all drylands ecosystems, Lal (2001) estimated that they had the potential to sequester up to 0.4–0.6 Gt of carbon a year if eroded and degraded dryland soils were restored and their further degradation were stopped. In addition, he suggested that various active ecosystem management techniques, such as reclamation of saline soils, could increase carbon sequestration by 0.5–1.3 Gt of carbon a year. Squires et al (1995) estimated similar figures. Keller and Goldstein (1998) reached the slightly higher figure of 0.8 Gt of carbon per year using estimates of areas of land suitable for restoration in woodlands, grasslands, and deserts, combined with estimates of the rate at which restoration can proceed.

Other studies have examined specific ecosystems in particular locations. For example, Glenday (2008) measured forest carbon densities of 58 to 94 tonnes C/ha in the dry Arabuko-Sokoke Forest in Kenya and concluded that improved management of wood harvesting and rehabilitation forest could substantially increase terrestrial carbon sequestration. Farage et al. (2007) used soil organic matter models to explore the effects of modifying agricultural practices to increase soil carbon stocks in dryland farming systems in Nigeria, Sudan and Argentina. Modelling showed that it would be possible to change current farming systems to convert these soils from carbon sources to net sinks without increasing farmers' energy demand. The models indicated that annual rates of carbon sequestration of 0.08-0.17 tonnes per ha per year averaged over the next 50 years could be obtained.

Despite these studies, significant gaps in knowledge remain. Better information is needed on the impact of land use changes and desertification on carbon sequestration and the cost-benefit ratio of soil improvement and carbon sequestration practices for small landholders and subsistence farmers in dryland ecosystems (MA 2005).

### **Linking drylands development and carbon markets**

There are two markets for carbon sequestration: a) the compliance market governed by the United Nations Framework Convention on Climate Change (UNFCCC) through its Kyoto Protocol and b) the voluntary market. The role of the natural biosphere in climate change mitigation is recognised in the UNFCCC through Land Use Land Use Change and Forestry (LULUCF).

Annex I Parties, under Article 3.3 of the Kyoto Protocol, can use "*direct human-induced land-use change and forestry activities, limited to afforestation, reforestation and deforestation since 1990, measured as verifiable changes in carbon stocks,*" to meet emissions reductions targets. In addition, they can elect Forest Management, Grassland Management, Cropland Management, and Revegetation for inclusion in the accounting process. There are calls by some to include all lands and associated processes in the LULUCF, rather than the narrow activities specified above.

The rules for LULUCF were only set after emission reduction targets had been agreed. This has been viewed as a limitation, as in effect land use activities ‘offset’ emissions in other sectors, rather than acting as an integral part of the mitigation portfolio. Issues still remain over the permanence of sequestration activities as management changes or natural disturbances can quickly release any carbon accumulated.

The opportunities for Non Annex 1 countries to participate in such activities is also limited, and restricted to the Clean Development Mechanism (CDM); where Annex I countries can gain carbon credits through activities in developing countries. CDM activities are restricted to Afforestation, Reforestation and Deforestation activities, and can make up only 1% of the emissions reduction portfolio for Annex I countries.

As yet few forestry-based carbon sequestration activities have been funded through the CDM, partly because of concerns about additionality, permanence and leakage. Voluntary markets have developed their own regulations and protocols, and are the only outlet for reduced deforestation programmes at the moment.

However, the UNFCCC is considering introducing a financial mechanism to reduce emissions from deforestation and forest degradation (REDD) in developing countries. There is still a great deal of uncertainty about the form of the mechanism, not least how it will be funded. One option is to do so through a specific fund, another is a market-based mechanism that would allow developing countries to sell carbon credits on the basis of successful reductions in emissions from deforestation and forest degradation. A market-based mechanism is expected to generate a much greater supply of funds; one estimate, based on a relatively low carbon price of U.S. \$10 per ton and an estimate of individual countries’ ability to slow deforestation, suggests a potential market of U.S. \$1.2 billion a year (Niles et al., 2002).

The United Nations Collaborative Programme on Reducing Emissions from Deforestation and Forest Degradation in Developing Countries (UN-REDD Programme) is a collaboration between FAO, UNDP and UNEP. It is aimed at “tipping the economic balance in favour of sustainable management of forests so that their formidable economic, environmental and social goods and services benefit countries, communities and forest users while also contributing to important reductions in greenhouse gas emissions”. Its immediate goal is to assess whether carefully structured payment structures and capacity support can create the incentives

预览已结束，完整报告链接和二维码如下：

[https://www.yunbaogao.cn/report/index/report?reportId=5\\_10661](https://www.yunbaogao.cn/report/index/report?reportId=5_10661)

