

# Prevention, identification and management of health worker infection in the context of COVID-19

Interim guidance

30 October 2020



## Key points

- Health workers in contact with and/or who care for COVID-19 patients are at a higher risk of infection than the general population. Mitigating and reducing this risk is essential to protecting their well-being and reducing the spread of COVID-19.
- Available scientific evidence suggests that appropriate personal protective equipment use, hand hygiene best practices, implementation of universal masking policies in health care facilities and adequate infection prevention and control (IPC) training and education are associated with decreased risk of COVID-19 among health workers.
- The prevention of SARS-CoV-2 infections in health workers requires a multi-pronged integrated approach that includes occupational health and safety (OHS) measures as well as IPC. All health-care facilities should establish or strengthen and implement (a) IPC programmes and (b) Occupational Health and Safety programmes with protocols to ensure HW safety and prevent HW infections while in the work environment. Ensuring adequate clinical staffing levels is recommended to prevent the transmission of health care-associated infections.
- Early detection of SARS-CoV-2 infection among health workers can be achieved through syndromic surveillance and/or laboratory testing and is a key strategy to prevent secondary transmission from health workers to patients, between health workers throughout health-care settings and from health workers to contacts outside of health facilities. A national and/or local surveillance and testing strategy should be developed and implemented.
- A system for managing exposures based on risk assessment should be in place to promote and support health workers' reporting of occupational and non-occupational exposures to or symptoms of COVID-19.
- A system for managing suspected infections, including measures for health workers who test positive for SARS-CoV-2 and those who are

symptomatic and test negative for SARS-CoV-2, should be in place.

- Clear criteria for returning to work should be established according to the WHO principles for discontinuing isolation for COVID-19.
- Health systems and facilities should maintain a blame-free culture with regards to COVID-19 infections in health workers.
- WHO has provided several tools for surveillance and studies to better understand the extent of infections and risk factors for SARS-CoV-2 infection among health workers.

## Background

Health workers,<sup>1</sup> in particular those in contact with and/or who care for COVID-19 patients, are at higher risk of being infected with SARS-CoV-2 than the general population.(1,2) Data collected by the World Health Organization (WHO) global surveillance for COVID-19, primarily from European and American countries, estimate that approximately 14% of COVID-19 cases reported to WHO are among health workers. Transmission of the SARS-CoV-2 virus to health workers has been documented to occur in both acute care and long-term care settings; from patients and residents to health workers as well as among health workers, also potentially associated with exposures to infected co-workers in common areas and break rooms.(3–7)

As the pandemic evolves, studies indicate that transmission involving health workers is also occurring in community settings (such as in households) in addition to health care settings.(6,8–12) COVID-19 infections among health workers may lead to a depleted workforce during a time when the demand on the health care system has increased. In addition, health workers who are infected are at risk of transmitting SARS-CoV-2 virus to others in households and other community settings. For more information on evidence of the epidemiology and risk factors of health worker infections see Box 1. An understanding of the transmission of SARS-CoV-2, as described in the WHO [Transmission of SARS-CoV-2: implications for infection prevention precautions\(1\)](#) a key element to implementing appropriate infection prevention measures.

<sup>1</sup> Health workers are defined by WHO as all people engaged in actions with the primary intent of enhancing health, including

social care workers who often have roles in the provision of care in long-term care facilities and in community settings. (61)

This document offers guidance for the prevention, surveillance and testing of COVID-19 in health workers<sup>1</sup> including the management following exposure and eventual safe return to work of health workers who have had suspected or confirmed SARS-CoV-2 infections. In the WHO surveillance database, the term health worker includes physicians, nurses, allied health workers (x-ray, laboratory staff, physiotherapists etc.), and administrative and support staff, such as cleaning and laundry personnel, admission/reception clerks, patient transporters and catering staff.(13)

Prevention of infection in the workplace requires a multi-pronged, integrated approach that includes IPC and occupational health and safety (OHS) measures plus adherence to public health and social measures in the community. Syndromic surveillance is a process that is often used by public health surveillance systems for early outbreak detection and focuses on early symptom identification.(14) Since early in the COVID-19 pandemic, laboratories have been using nucleic acid amplification tests (NAATs), such as real time reverse transcription polymerase chain reaction (rRT-PCR) assays, to detect SARS-CoV-2, the virus that causes COVID-19 disease. Antigen-detecting tests (Ag-RDTs) are now entering the armamentarium of tools that can play a significant role in guiding patient management, public health decision making and surveillance of COVID-19.(15)

This guidance document complements the WHO [Risk assessment questionnaire](#).(16) on management of health workers' infections in the context of COVID-19 and [Coronavirus disease \(COVID-19\) outbreak: rights, roles and responsibilities of health workers, including key considerations for occupational safety and health: interim guidance](#).(17) This guidance is intended for both the national and facility levels, for public health authorities, health facility administrators, occupational health departments and infection prevention and control (IPC) departments or focal points<sup>2</sup> and may be adapted according to national and local contexts.

Rapid literature reviews were conducted to gather the evidence basis for the development of this document, in particular regarding epidemiology of and risk factors for SARS-CoV-2 infections among health workers and on effectiveness of active and passive syndromic surveillance and routine laboratory testing. This guidance document has been developed by consulting the WHO ad-hoc COVID-19 IPC Guidance Development Group (COVID-19 IPC GDG), as well as external experts and WHO staff with expertise in the field of occupational health and health work forces' rights and development. These groups reviewed the available evidence, considered various country experiences and formulated this consensus-based advice for syndromic surveillance and testing of health workers.

### **Box 1: Evidence on the epidemiology of and risk factors for health worker infection**

Limited data are available globally and at country level about SARS-CoV-2 infection among health workers. WHO has recently produced a summary of health worker SARS-CoV-2 infections.(18) Briefly, WHO global surveillance for COVID-19, primarily in countries in the WHO European and American Regions, indicate that approximately 14% of COVID-19 cases reported to WHO are identified as occurring in health workers. Even among countries with >75% completion of variables for the data submitted related to health worker status, the proportions of health workers infected varied broadly, ranging from 2-35%. Timing of reporting, fluctuation of community transmission patterns and facility implementation of IPC measures had an impact on occurrence of health workers' infections. A recent report from the International Council of Nurses, who conducted a survey of 50 countries, mostly from the Europe and the Americas, found that health worker infections ranged from 1-32% of all confirmed COVID-19 cases.(19)

Limited availability of published data on health worker infections can be attributed in part to difficulties in distinguishing infection acquisition in community outbreaks from that in health-care settings and differences in data confidentiality aspects of health worker surveillance. In general, there are limitations to obtaining comprehensive data, such as high variability in the completeness of the information.

Studies cited in a living rapid review commissioned by WHO on the epidemiology and risk factors for COVID-19 and other coronaviruses (SARS-CoV-1 and MERS-CoV), in health workers,(3) found that estimates of SARS-CoV-2 infections among health workers vary significantly across studies. The incidence of SARS-CoV-2 infection (PCR-positive) ranged from 0.4% to 49.6%, and the prevalence of SARS-CoV-2 seropositivity ranged from 1.6% to 31.6%, depending on the study. Factors contributing to increased risk of occupationally acquired SARS-CoV-1, MERS CoV, or SARS-CoV-2 infection include variability in estimates include differences in settings, type and length of exposure(s), increased intensity of community transmission where health facilities are, presence and severity of symptoms and inadequate use and supplies of personal protective equipment (PPE), among others. Some key findings of the living review include (3):

- Appropriate personal protective equipment use, hand hygiene best practices, implementation of universal masking policies in health-care facilities, and adequate infection prevention and control (IPC) training and education for all HWs are associated with decreased risk of infection in health workers.
- Available evidence does not find an association between age and sex or health worker role (e.g. nurse versus physician) and risk of SARS-CoV-2 infection.
- SARS-CoV-2 infections have been observed in various hospital departments and health workers performing in different roles, including those without direct patient contact.
- Certain exposures (e.g., performing intubations, direct patient contact, and contact with bodily secretions) and inconsistent/incomplete use of PPE are associated with increased risk of coronavirus infections in health workers.

<sup>2</sup> IPC focal point is defined as a professional appointed to be in charge of IPC at the national, sub-national or facility/organization level.(62)

Transmission of the SARS-CoV-2 virus to health workers has been documented in acute care and long-term care settings: from patients and residents to health workers from one health worker to another, including possible exposures in common areas and break rooms.(3–7) Seroprevalence and genomic studies have been and are currently being conducted among health workers. Studies indicate that transmission involving health workers also occurs in community settings (such as households) in addition to health care settings.(6,8–12)

While appropriate use of PPE is one critical protective measure for health workers, strategies to mitigate harms associated with prolonged and reuse of PPE and other identified risk factors are described in the WHO interim guidance: [Coronavirus disease \(COVID-19\) outbreak: rights, roles and responsibilities of health workers, including key considerations for occupational safety and health: interim guidance.](#) (17)

## Key principles for preventing infections in health workers

The prevention of SARS-CoV-2 infections in health workers requires a multi-pronged integrated approach of IPC and occupational health and safety (OHS) measures.(17,20) WHO recommends that all health care facilities should establish and implement IPC programmes and OHS programmes with protocols to ensure health worker safety and prevent infections with COVID-19 in the work environment.(20)

Studies report that health workers in areas impacted by COVID-19 experience high levels of depression, anxiety, and psychological distress.(21–23) Health worker shortages and long shifts without adequate rest periods and shortages of personal protective equipment are important determinants leading to fatigue, and inadequate adherence to infection prevention practices.(21,22) This has been highlighted by WHO in guidelines, in which adequate staffing levels and adequate IPC training are strongly recommended as a core components of effective IPC programmes to prevent health care-associated infections, including those spread through outbreaks.(20) Lack of adequate social health protection measures such as health monitoring, sick leave for quarantine and “stay home if unwell” policies for certain groups of health workers (e.g. self-employed private providers, community health workers and lay providers) has also been reported. Therefore, WHO has previously recommended the establishment of national IPC and OHS programmes at the national level and in all health care facilities.(24)

Below is a list of existing and new recommendations to prevent SARS-CoV-2 health worker infections.

### 1. Establish an infection prevention and control programme

The [WHO Guidelines on core components of infection prevention and control programmes at national and acute health care facility levels](#) (20) are the foundation of WHO strategies to prevent current and future threats from infection and antimicrobial resistance in health care. A facility level IPC programme with a dedicated and trained IPC team, or at minimum an IPC focal point, should be in place and supported by the national and facility senior management.(20) Ensuring adequate clinical staffing levels is recommended as a core component to prevent the transmission of health care-associated infections, in particular spread through outbreaks. [Minimum requirements](#)(25) have been identified to facilitate the step-wise implementation of the WHO core components for IPC programmes, in particular in countries where IPC is limited or non-existent.(25) Achieving the IPC minimum requirements and more robust and comprehensive IPC programmes according to all WHO IPC core components across the whole health system in all countries, is essential to sustaining efforts to

control the COVID-19 pandemic and other emerging infectious diseases and prevent health care-associated infections and antimicrobial resistance.(20,25)

Long-term care services have been identified as high risk for transmission of COVID-19 between residents and staff.(7,26) WHO guidance specific to these settings has been developed: [Preventing and managing COVID-19 across long-term care services](#); policy brief,(27) [Preventing and managing COVID-19 across long-term care services](#) web annex(28) and [Infection prevention and control guidance for long-term care facilities in the context of COVID-19](#).(29) They should be utilized in addition to the above IPC documents.

Specific IPC measures, recommended by WHO, to reduce transmission of SARS-CoV-2 among health workers are described in a number of key IPC technical guidance documents,(2,29–33) and include the following:

- Ensure triage, early recognition, and source control (isolating suspected and confirmed COVID-19 cases including long-term care residents,
- Apply standard IPC precautions for all patients, with special attention to appropriate hand hygiene and environmental cleaning,
- Implement additional precautions (droplet and contact and, wherever applicable, for aerosol-generating procedures, airborne precautions) for suspected and confirmed cases of COVID-19 universal medical masking by health workers in health-care facilities, including in common areas where health workers interact,
- Implement administrative controls, such as IPC policies and procedures, including appropriate behaviours and compliance with key IPC measures in common areas,
- Use or introduce environmental and engineering controls such as appropriate ventilation.

### 2. Establish an occupational health and safety programme

All health services should have an occupational health and safety policy and programme including occupational health focal point or occupational health service; labour-management committee for health and safety; regular workplace risk assessment covering all hazards and the effectiveness of their controls; immunizations; blame-free reporting of accidental/unprotected exposures to pathogens and incidents; medical surveillance, education and training of workers; and hygiene measures.(34) Specific measures to protect health workers from occupational risks amplified by the COVID-19 pandemic are described in the WHO interim guidance “COVID-19: Occupational health and safety of health workers, right and responsibilities” (forthcoming)” and WHO/ILO [“Occupational safety and health in public health](#)

[emergencies: a manual for protecting health workers and responders](#)".(35)

A key element for transmission prevention and control in health care settings is the application of engineering, environmental and administrative controls in addition to individual behaviours and PPE. In addition to the core elements of IPC and OHS programmes described above, the following measures should be included to prevent health worker infections:

- Regular assessment of risks and effectiveness of control measures, including compliance with IPC and safety protocols and occupational risk assessment,
- Education and training of all staff on IPC measures and occupational health and safety, including regular refresher training,
- Access to and appropriate use of supplies for IPC such as hand hygiene supplies and PPE (medical masks, respirators, eye protection, gloves, gowns), that should be available in sufficient quantity and size ranges, and meeting quality standards,
- Monitoring of IPC procedures and regular feedback to various audiences including clinical staff, supported by mentoring and supervision for practice; and reinforcement of skills to establish strong social norms related to IPC adherence,(36)
- Monitoring behavioural and social barriers and enablers for health worker adherence, such as perceptions about the value of procedures, confidence about following procedures and perceptions about available support, (36)
- Occupational health and safety policies and procedures including:
  - staff screening and testing, staff illness protocols and safe return-to-work policies
  - policies to allow staff to stay home if unwell, without loss of income
  - procedures for blame-free reporting and investigation of unprotected exposures and contacts with suspected or confirmed COVID-19 cases
  - management protocols to ensure sufficient staff; safe staff-to-patient ratios; appropriate shifts; rest periods in areas with adequate space and ventilation; and reminders to staff to continue adherence to IPC procedures
- Regular communication between staff and senior management, including staff participation in planning,
- Cooperation between employers and sub-contractors operating in the same health facility in developing and implementing safety protocols and protective measures.

### Early detection of SARS-CoV-2 infections in health workers to prevent further transmission

Early detection of COVID-19 infection among health workers can be achieved through syndromic surveillance and/or laboratory testing and is a key strategy to prevent secondary transmission to patients, between health workers and throughout health-care settings.

Syndromic surveillance can be conducted using passive methods (e.g. relying on self-reporting of symptoms or illness by HWs) or active methods (e.g. that involves interviewing or assessing HWs to identify suspected cases of the disease under surveillance).(14,37)

Fever is a common symptom of COVID-19. A systematic review found that fever, myalgia or arthralgia, fatigue, and headache are common among COVID-19 patients.(38,39) Loss of taste (ageusia) and smell (anosmia), ocular pain, general malaise, and extreme tiredness have also been reported.(6,7) Some cases have not reported any symptoms.(40)

Limited available studies have found that PCR testing is more frequently positive among symptomatic health workers compared to those without symptoms (odds ratio ranging from 3.5-19.4).(12,41–44)The proportion of health workers who tested positive while asymptomatic has been found to range between 12- 23.1%.(11,26,41,43,44)

A small study from Scotland, United Kingdom, in which health workers who reported symptoms were promptly tested for COVID-19, rather than quarantined for up to 14 days, noted that testing may have saved the health system approximately 8573 lost workdays due to reduced staff absences.(45) A recent large multicenter study of long-term care facilities conducted in the United States of America found that 1.3 cases of COVID-19 were found in health workers for every 3 cases identified among residents. This finding is consistent with other studies conducted in long-term care settings that found cases of COVID-19 in health workers when a wide testing strategy for all health workers was implemented once a resident was identified as positive for COVID-19(7).

In general, while studies show that testing health workers on a regular schedule is likely to identify infection, clear intervals for routine testing or time points have not been identified.(46–49)

Based on available evidence, WHO advises the following:

#### **1. Syndromic surveillance of health workers for COVID-19 symptoms should be performed before they enter the workplace.** This should include:

- Passive surveillance: encourage health workers to report symptoms to the occupational health professional or another designated officer in the facility before their shift (including via routine digital reporting forms where available), and during or after their shift.
- Active surveillance: establish a confidential process for ensuring health workers are screened for symptoms of COVID-19, including fever, and any potential exposure risks on arrival for their shift.

Passive surveillance may be the only option when resources are limited, but active syndromic surveillance should be considered if human resources and logistics permit it. All efforts to institute active syndromic surveillance are recommended when there are clusters of transmission in the health facility or in the areas where the health facility is

located. If there is community transmission<sup>3</sup>, syndromic surveillance is critical.

Symptoms to be monitored for syndromic surveillance of health workers should include, at minimum: fever, dry cough, myalgia, arthralgia, fatigue, headache, shortness of breath, anosmia and ageusia. Staff with any of the above symptoms

or who fail the screening process should contact their OHS for further instructions. Employment policies should be in place, such as sick leave and ability to stay home if unwell, that grant confidentiality and are non-punitive for health workers who become contacts<sup>4</sup>, or infected with SARS-CoV-2.(17)

**Table 1. Examples of syndromic surveillance approaches**

COVID-19 Transmission scenario (50)	Type of syndromic surveillance for health workers	Possible approach
No cases or Sporadic cases	Implement passive syndromic surveillance	<ul style="list-style-type: none"> <li>Staff self-report to occupational health or other designated officer if they experience any symptoms including fever.</li> </ul>
Clusters of cases	Implement passive syndromic surveillance, consider active surveillance if resources available	<ul style="list-style-type: none"> <li>Staff self-report to occupational health or other designated officer if they experience any symptoms including fever.</li> <li>If resources are available, consider a process to actively monitor staff for symptoms including fever.</li> </ul>
Community transmission	Implement active syndromic surveillance	<ul style="list-style-type: none"> <li>A process is put in place in which staff temperatures are monitored and staff are assessed actively (screened) for symptoms at the beginning of each shift at minimum.</li> </ul>

## 2. National and sub-national testing strategies for health workers for detection of SARS-CoV-2 infections should be developed and implemented.

Adequate laboratory testing for SARS-CoV-2 infections is another element needed to more accurately identify SARS-CoV-2 transmission among health workers.(15,51) When considering a testing strategy, the following contextual factors should be taken into account: effectiveness of the facility's occupational health and IPC programmes (including protocols fully implemented by employers/management and demonstrated staff adherence to protocols), the local transmission scenario, available resources and infrastructures for testing, and the impact of COVID-19 on the health workforce (e.g. potential absences due to sick leave, self-isolation or quarantine). In settings with limited resources in areas of community transmission, WHO recommends that health workers be prioritized for testing, regardless of whether they are a contact of a confirmed case (to protect health workers and reduce the risk of nosocomial transmission).(52)

WHO provides recommendations for RT-PCR and antigen-based testing for the diagnosis of SARS CoV-2.(15,51)

The testing strategy should include:

### a) Testing health workers following exposure to SARS-CoV-2

Health workers in a health-care facility who are contacts<sup>4</sup> of a suspected or confirmed case(53), as a result of an

unprotected exposure at work or in the community, should consult the occupational health focal point to be assessed using the [WHO risk assessment and management of exposure of health care workers tool](#).(16) WHO recommends that all contacts with high-risk exposure should be tested for SARS-CoV-2.

### b) Routine testing of health workers for COVID-19 surveillance

The need for routine testing should be decided using a risk-based approach and the following factors should be taken into account:

- The intensity of transmission in the setting of the health facility(ies), for example in the presence of community transmission or intense outbreaks of COVID-19.
- The capacities of the facility and laboratories to conduct the testing including financial and human resources available, as well as availability of testing materials and laboratory capacity.
- The volume of patients identified as positive for SARS-CoV-2, admitted to the facility or being assessed by health workers.
- The positivity rate among staff.
- The number of staff who are ill but not diagnosed with COVID-19 and in quarantine as contacts for COVID-19, leading to inability to provide adequate and safe staffing levels.

<sup>3</sup> Community transmission is described as outbreaks with the inability to relate confirmed cases through chains of transmission for a large number of cases, or by increasing positive tests through sentinel samples(50)

<sup>4</sup> A contact is a person who has experienced any one of the following unprotected exposures during the 2 days before and the

14 days after the onset of symptoms of a probable or confirmed case: 1. face-to-face contact with a probable or confirmed case within 1 metre and for at least 15 minutes 2. direct physical contact with a probable or confirmed case 3. direct care for a patient with probable or confirmed COVID-19 disease without using recommended personal protective equipment OR 4. other situations as indicated by local risk assessments(53)

**c) Testing health workers in long-term care facilities**

Irrespective of the COVID-19 transmission scenario, health workers staffing or working with long-term care facilities should be considered for routine testing, and at a

minimum be tested for COVID-19 as soon as a positive case of COVID-19 is identified in either residents or staff.

Table 2 provides some examples of scenarios for the application of a risk-based approach.

**Table 2: Examples of scenarios and testing strategies for health workers**

Health-care Setting	Transmission scenario(50)	Possible testing strategy target to consider (where resources allow)
Acute care	No cases or Sporadic cases	<ul style="list-style-type: none"> <li>• Symptomatic health workers</li> <li>• Health worker identified as a contact of a SARS-CoV-2 case               <ul style="list-style-type: none"> <li>– Health workers associated with transmission to or from a patient or resident or with an outbreak investigation</li> </ul> </li> </ul>
	Clusters or Community transmission	<ul style="list-style-type: none"> <li>• Symptomatic health workers</li> <li>• Health worker identified as a contact of a SARS-CoV-2 case               <ul style="list-style-type: none"> <li>– Health workers associated with transmission to or from a patient, a cluster, or with an outbreak investigation</li> </ul> </li> <li>• Health workers working in any clinical area, identifying priority areas based on risk assessment (e.g. triage, emergency services or COVID-19 wards) where resources are limited</li> <li>• All health workers who work in COVID-19 services or facilities</li> </ul>
Long-term care	All transmission scenarios	<ul style="list-style-type: none"> <li>• Symptomatic health workers</li> <li>• Health workers identified as a contact of a SARS-CoV-2 case</li> <li>• Testing of all health workers when a positive case of SARS-CoV-2 is identified in a resident or staff member</li> <li>• Routine testing of health workers, if feasible</li> </ul>

The frequency of health worker testing will depend on the level of transmission within a facility and surrounding areas, objective(s) of the testing strategy (i.e. surveillance versus outbreak control), capacity of the facility and relevant laboratories to conduct the testing and national and local guidance. During an outbreak of COVID-19, testing should be conducted regularly (e.g. weekly, if resources allow) until there are no cases of COVID-19 in health workers or residents in the facility.(7,26,48)

### Managing health worker exposures, infections and safe return to work

A blame-free system for managing health worker exposures to COVID-19 should be in place to promote and support reporting of exposures or symptoms. Organizations providing health care should have paid sick leave policies for health workers that are non-punitive, not associated with any financial disincentives, confidential, flexible and consistent with public health guidance. OHS focal points should keep confidential records of health workers who are exposed to COVID-19 and monitor those who develop or report symptoms or test positive.

A key element for developing policies for managing health worker exposures, infections and the safe return to work is available evidence about the duration of viral shedding among COVID-19 patients and in particular the timeframe within which replication-competent virus can be isolated. The WHO scientific brief [Criteria for releasing COVID-19](#)

[patients from isolation\(54\)](#) provides an overview of the evidence.

Accordingly, the following guidance is based on several studies showing that in patients with mild to moderate COVID-19, replication-competent virus has not been recovered after 10 days following symptom onset.

WHO provides the following advice:

#### **1. Health workers should be encouraged to report both occupational and non-occupational exposures to COVID-19.**

If a health worker reports an unprotected exposure to COVID-19, fails syndromic screening on arrival or develops symptoms during their shift, clear policies and procedures should be in place outlining the steps that should be taken, which include:

- Instructions for the health worker to immediately stop working, put on a medical mask if not already wearing one, report to their OHS officer and self-isolate.
- OHS should meet with the health worker to conduct an assessment and exposure history where resources permit, or ask the health worker to complete and submit the form for the WHO Risk assessment and management of exposure of health care workers in the context of COVID-19.
- OHS should identify a risk categorization based on the risk assessment tool for a health worker who has had an unprotected exposure and determine appropriate management, including the health worker's ability to return to work.

- OHS should contact local public health authorities to notify them about health workers who report both occupational- and non-occupational related exposures and to arrange appropriate follow-up and monitoring.
- The occupational disease should be reported according to OHS Acts.
- Strategies to mitigate workforce shortages should be in place.(17,34,55,56)

Further information on risk assessment and management of COVID-19 exposures among health workers can be found [here](#).(16) The approach indicated in this guidance distinguishes exposures with high and low risk for COVID-

19 infection. The key advice for different situations is summarized in Table 3.

Based on risk classification post-exposure, the occupational health and safety department can advise the health worker to:

- Continue to work depending on ability to do so and the exposure risk assessment,
- Provide recommendations to monitor symptoms and for additional follow-up as needed,
- Arrange for testing for SARS-CoV-2 according to the national and local testing strategy,
- Consider quarantine, depending on the nature of the exposure.

**Table 3: Health worker exposure risk and advised actions**

Exposure type	Health worker status	Advice
<b>Lower risk exposure</b> in the workplace: <ul style="list-style-type: none"> <li>• provided direct care to COVID -19 patient while wearing required PPE and following IPC precautions,</li> <li>• present during an AGP on a patient with COVID-19 while wearing required PPE and following IPC precautions,</li> <li>• exposure to colleague who is a suspect or COVID-19 positive case at work while wearing a mask.</li> </ul>	No symptoms (asymptomatic)	<ul style="list-style-type: none"> <li>• May continue to work following IPC measures including local requirements for wearing of masks.</li> <li>• Test for SARS-CoV-2, if resources available. Follow guidance for <a href="#">Diagnostic Testing for SARS CoV-2.(51)</a></li> <li>• Reinforce IPC measures (physical distancing, hand hygiene, PPE and use of masks.</li> <li>• Self-monitor for symptoms for 14 days and report immediately to OHS if any symptoms develop.</li> <li>• If positive, identify contacts and follow up according to contact tracing procedures.</li> </ul>
	Symptomatic	<ul style="list-style-type: none"> <li>• Staff member self isolates.</li> <li>• Monitor with OHS.</li> <li>• Test for SARS-CoV-2. Follow guidance for <a href="#">Diagnostic Testing for SARS CoV-2.(51)</a></li> <li>• If positive, identify contacts and follow up according to contact tracing procedures.</li> </ul>
<b>Higher risk exposure</b> in the workplace: <ul style="list-style-type: none"> <li>• provided direct care to COVID-19 patient with no or inappropriate PPE, or a breach in PPE integrity or other IPC precautions not followed (i.e. hand hygiene not performed as per the WHO 5 moments, lack of cleaning and disinfection of surface/environment),</li> <li>• present during an AGP without or inappropriate PPE, breach in PPE integrity, or other IPC precautions not followed (e.g. hand hygiene not performed per the WHO 5 moments, lack of cleaning and disinfection of surface/environment),</li> <li>• exposure (&gt;15 min face-face contact, &lt; 1m) to a colleague who is identified as positive for COVID-19 with no masks (e.g. in a break room, while eating etc.),</li> <li>• exposure to splash or spray of body fluids/blood and/or a puncture/sharp injury.</li> </ul>	No symptoms (asymptomatic)	<ul style="list-style-type: none"> <li>• Staff to quarantine for 14 days after last exposure.</li> <li>• Staff to remain off work for 14 days from last exposure.</li> <li>• Test for SARS CoV-2. Follow guidance for <a href="#">Diagnostic Testing for SARS CoV-2(51)</a></li> <li>• If positive, identify contacts and follow up according to contact tracing procedures.</li> <li>• Monitor daily for symptoms and notify OHS.</li> </ul>
	Symptomatic	<ul style="list-style-type: none"> <li>• Staff member self-isolates.</li> <li>• Test for SARS-CoV-2. Follow guidance for <a href="#">Diagnostic Testing for SARS-CoV-2.(51)</a></li> <li>• Identify contacts and follow up according to contact tracing procedures.</li> <li>• See guidance below for return to work.</li> </ul>

Non-occupational exposure (e.g. contact <sup>4</sup> with a confirmed cases who is a family or community member).	Asymptomatic	<ul style="list-style-type: none"> <li>Quarantine for 14 days after the last exposure.</li> <li>If positive, identify contacts and follow up according to contact tracing procedures.</li> </ul>
	Symptomatic	<ul style="list-style-type: none"> <li>Staff member to isolate.</li> <li>Test for SARS-CoV-2.</li> <li>Follow guidance for <a href="#">Diagnostic Testing for SARS-CoV-2</a>.</li> <li>If positive, identify contacts and follow up according to contact tracing procedures.</li> <li>See the guidance below for return to work.</li> </ul>

## 2. Managing health worker infections

Any health worker who identifies as symptomatic or tests positive for SARS-CoV-2 should:

- immediately be isolated and stop all patient care activities,
- inform their supervisor who should notify the IPC and OHS,
- seek care if feeling unwell or symptoms worsen through the appropriate referral system.

The following table outlines the advice and the management steps in the event a HW tests positive for SARS-CoV-2.

**Table 4: Measures for health workers positive for SARS-CoV-2**

Health worker status	IPC Measures
Health worker tests SARS-CoV-2 positive (with or without symptoms)	<ul style="list-style-type: none"> <li>Isolate in a health care facility, designated setting (e.g. health-care facility, non-traditional facility), or at home (51) as appropriate and according to clinical condition for a minimum of 10 days plus 3 days without symptoms (33)</li> </ul>
Health worker is symptomatic but tests negative for SARS-CoV-2	<ul style="list-style-type: none"> <li>Follow guidance for <a href="#">Diagnostic Testing for SARS-CoV-2</a>,(51)</li> <li>Consult with the OHS on whether to return to work and consider if additional testing is required for alternate diagnoses according to local guidance,</li> <li>Any health worker permitted to return to work should be advised of symptoms to monitor and follow infection control guidance as described above, including the use of appropriate PPE.</li> </ul>

If a health worker's infection is related to an occupational exposure such as incorrect IPC practices, appropriate corrective measures, such as refresher training for staff on IPC measures, should be put in place to address and correct breaches. Facilities should ensure adequate supplies of appropriately fitted PPE are available for health workers and that processes are in place for monitoring and observation of IPC procedures, including fit checking of respirators, and correct order for PPE removal and disposal. There also should be workplace reminders to use hygiene measures during work

criteria for releasing COVID-19 patients from isolation are as follows:(54)

- Symptomatic patients may be released from isolation 10 days after symptom onset, plus at least 3 additional days without (including without fever<sup>5</sup> and without respiratory symptoms).
- Asymptomatic individuals can be released from isolation 10 days after they first tested positive.

Some individuals may experience symptoms (such as a nose

预览已结束，完整报告链接和二维码如下：

[https://www.yunbaogao.cn/report/index/report?reportId=5\\_24306](https://www.yunbaogao.cn/report/index/report?reportId=5_24306)

