

**CARING FOR NEWBORNS AND
CHILDREN IN THE COMMUNITY**

**A TRAINING COURSE FOR
COMMUNITY HEALTH WORKERS**

Caring for the sick child in the community

Adaptation for high HIV or TB settings



CHART BOOKLET



Caring for the sick child in the community: a training course for community health workers, adaptation for high HIV or TB settings. Chart booklet
ISBN 978-92-4-001736-8 (electronic version)
ISBN 978-92-4-001737-5 (print version)

This publication was originally published in 2014.

© World Health Organization 2020

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>). Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: “This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition”. Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<http://www.wipo.int/amc/en/mediation/rules/>).

Suggested citation. Caring for the sick child in the community: a training course for community health workers, adaptation for high HIV or TB settings. Chart booklet. Geneva: World Health Organization; 2020. Licence: [CC BY-NC-SA 3.0 IGO](#).

Cataloguing-in-Publication (CIP) data. CIP data are available at <http://apps.who.int/iris>.

Sales, rights and licensing. To purchase WHO publications, see <http://apps.who.int/bookorders>. To submit requests for commercial use and queries on rights and licensing, see <http://www.who.int/about/licensing>.


Third-party materials. If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement. The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters. All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

Cover photo J. Lucas

OVERVIEW: CARING FOR THE SICK CHILD IN THE COMMUNITY

(child age 2 months up to 5 years)





IDENTIFY PROBLEMS: ASK AND LOOK

ASK the caregiver: What are the child's problems?


LOOK at the child.

- D Cough? If yes, for how long? _ days
- D Diarrhoea (3 or more loose stools in last 24 hours)? If yes, for how long? _ days.
- D If diarrhoea, blood in stool?
- D Fever (reported or now)? If yes, started _ days ago.
- D Convulsions?
- D Difficulty drinking or feeding? If yes, not able to drink or feed anything?
- D Vomiting? If yes, vomits everything?
- D Has HIV?
- D At risk of HIV because
 - D One or both parents have HIV and child has not tested for HIV? or
 - D Parents' current HIV status is unknown?
- D Lives in household with someone on TB treatment?
- D Any other problem?

- D Chest indrawing?
- D If cough, count breaths in 1 minute: _ breaths per minute (bpm).
- D Unusually sleepy or unconscious?
- D For child age 6 months up to 5 years, MUAC strap colour: _
- D Swelling of both feet?



DANGER SIGNS



--+ IF ANY DANGER SIGN, REFER CHILD URGENTLY TO HEALTH FACILITY

- D Cough for 14 days or more
- D Diarrhoea for 14 days or more
- D Blood in stool
- D Fever for last 7 days or more
- D Convulsions
- D Not able to drink or feed anything
- D Vomits everything
- D Has HIV and any other illness
- D Chest indrawing
- D Unusually sleepy or unconscious
- D For child age 6 months up to 5 years
 - D Red on MUAC strap, or
 - D Yellow on MUAC strap and has HIV
- D Swelling of both feet

--+ Assist referral to health facility:

- + Explain why child needs to go to the health facility.
- + GIVE FIRST DOSE OF TREATMENT:
 - + If diarrhoea, and if child can drink, begin giving ORS solution right away.
 - + If fever AND: convulsions; or unusually sleepy/ unconscious; or not able to drink or feed; or vomits everything, give rectal artesunate suppository (100 mg):
 - Age 2 months up to 3 years—1 suppository
 - Age 3 years up to 5 years—2 suppositories
 - + If fever AND danger sign other than the 3 above, give first dose of oral antimalarial AL:
 - Age 2 months up to 3 years—1 tablet
 - Age 3 years up to 5 years—2 tablets
 - + If fast breathing or chest indrawing, give first dose of oral antibiotic (amoxycillin tablet—250 mg):
 - Age 2 months up to 12 months—1 tablet
 - Age 12 months up to 5 years—2 tablets




- + For any sick child who can drink, advise to give fluids and continue feeding.
 - + Advise to keep child warm, if child is NOT hot with fever.
 - + Write a referral note.
 - + Arrange transportation, and help solve other difficulties in referral.
- FOLLOW UP child on return at least once a week until child is well.

To give ORS solution, see page 9.

To give first dose oral medication, see page 10

-+ SICK BUT NO DANGER SIGN?

- D Cough (less than 14 days)
- D Diarrhoea (less than 14 days AND no blood in stool)
- D Fever (less than 7 days) in a malaria area
- D At risk of HIV because
 - D One or both parents have HIV and child has not tested for HIV, or
 - D Parents' current HIV status is unknown
- D Lives in household with someone on TB treatment
- D Fast breathing:
 - In a child age 2 months up to 12 months, 50 breaths or more per minute
 - In a child age 12 months up to 5 years, 40 breaths or more per minute
- D Yellow on MUAC strap (does not have HIV)



If SICK but
NO danger
sign

TREAT at home
and
ADVISE
caregiver

Go to next page

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_24501

