



Rapid Communication:
Key changes to treatment of multidrug- and
rifampicin-resistant tuberculosis
(MDR/RR-TB)

August 2018

© World Health Organization 2018

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence.

If you create a translation of this work, you should add the following disclaimer along with the suggested citation: "This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition".

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization.

Suggested citation. Rapid communication: key changes to treatment of multidrug- and rifampicin-resistant tuberculosis (MDR/RR-TB). Licence: CC BY-NC-SA 3.0 IGO.

Cataloguing-in-Publication (CIP) data. CIP data are available at <http://apps.who.int/iris>.

Sales, rights and licensing. To purchase WHO publications, see <http://apps.who.int/bookorders>. To submit requests for commercial use and queries on rights and licensing, see <http://www.who.int/about/licensing>.

Third-party materials. If you wish to reuse material from this work that is attributed to a third-party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

Rapid Communication:

Key changes to treatment of multidrug- and rifampicin-resistant tuberculosis (MDR/RR-TB)

Background

Providing evidence-based guidelines to inform public health service delivery for Member States and other stakeholders is one of the core responsibilities of the World Health Organization (WHO).

The most recent WHO evidence-based guidelines for the treatment of multidrug-resistantⁱ or rifampicin-resistant tuberculosis (MDR/RR-TB) were published in October 2016. Subsequently, new evidence prompted a [public call for data](#) by WHO in anticipation of a formal review. Anonymized individual data received from clinical trials, cohort/observational studies and programmatic implementation of both longer and shorter MDR-TB regimens were incorporated into an individual MDR/RR-TB patient data base (IPD) hosted by McGill University, Canada, under contract with WHO.

[Cochrane](#) international methods for meta-analysis were followed to assess the relative contributions of individual medicines to patient treatment outcomes and to inform the design of MDR/RR-TB treatment regimens, the duration of treatment and the impact of drug resistance profiles on those outcomes.

WHO convened a Guideline Development Group meeting on 16-20 July 2018 to assess the results of these analyses using the international [GRADE](#) system (Grading of Recommendations, Assessment, Development and Evaluation for scientific evidence assessment and development of evidence-based policy guidelines and recommendations. The new MDR-TB treatment guidelines will be released later in 2018. These guidelines will replace all previous and current WHO guidelines on treatment of MDR/RR-TB.

This Rapid Communication aims to inform TB programme managers and other stakeholders of WHO Member States about the key implications for MDR-TB treatment regimens arising from the new evidence assessment. It highlights the immediate steps to be taken to ensure that MDR/RR-TB patients receive treatment in accordance with the latest evidence on effectiveness and safety.

While understanding that it would not be immediately possible to achieve the new standards of care in every individual MDR-TB patient, strategic planning should start immediately to enable rapid transition to the upcoming new WHO guidelines.

Data sources

- An IPD database with over 12,000 patient records from 50 studies of longer MDR-TB regimens;
- New data from 26 countries following the WHO public call, including the use of the shorter regimen in African and Asian settings and the use of bedaquiline worldwide;
- Aggregated results from the Otsuka phase III randomised controlled trial of delamanid, released in October 2017 and initially assessed by WHO in an expedited review in January 2018;

ⁱ Combined resistance to rifampicin and isoniazid, the two most important anti-TB drugs.

- Aggregated final results from the STREAM Stage 1 randomised controlled trial of a 9-month shorter MDR-TB regimen, after release of the interim results in October 2017 and initial assessment by WHO in an expedited review in February 2018.
- Pharmacokinetic and safety data from trials of bedaquiline and delamanid in children to consider the extension of treatment recommendations to children and adolescents.

Final results (outcomes) of treatment were used for developing policy recommendations. Patient data series that only reported interim results (such as 6-month culture conversion) were not considered, given that correlation between such surrogate endpoints and final outcomes (such as cure or treatment failure) has not been reliably established.

Treatment principles

- **Ahead of enrolment on MDR-TB treatment, all patients should receive appropriate counselling to enable informed and participatory decision-making.**
- **Patient information material needs to reflect the new changes so that patients are appropriately informed about their treatment options.**
- **Social support to enable adherence to treatment is very important to ensure a patient-centred approach to the delivery of care.**
- **Active TB drug safety monitoring and management (aDSM) is essential for all patients enrolled on MDR-TB treatment.**

Key medicine changes

Longer MDR-TB regimensⁱⁱ

The revised grouping of TB medicines recommended for use in longer MDR-TB regimens is presented in [Table 1](#). Medicines have been regrouped into **three categories and ranked** based on the latest evidence about the balance of effectiveness to safety:

- **Group A:** Medicines to be prioritised: levofloxacin/moxifloxacin, bedaquiline and linezolid
- **Group B:** Medicines to be added next: clofazimine, cycloserine/terizidone
- **Group C:** Medicines to be included to complete the regimens and when agents from Groups A and B cannot be used: ethambutol, delamanidⁱⁱⁱ, pyrazinamide, imipenem-cilastatin, meropenem, amikacin (streptomycin), ethionamide/prothionamide, *p*-aminosalicylic acid;

Medicines no longer recommended are kanamycin and capreomycin, given increased risk of treatment failure and relapse associated with their use in longer MDR-TB regimens. Use of amikacin did not show a

ⁱⁱ Longer MDR-TB regimens usually last 18-20 months and may be standardized or individualized. These regimens are usually designed to include at least five medicines considered to be effective.

ⁱⁱⁱ The position of delamanid will be re-assessed once individual patient data from Otsuka trial 213 has been reviewed; these data were not available for the evidence assessment outlined above.

similar association, although the same safety concerns as for the other injectables apply. Amoxicillin-clavulanic acid is only to be used to accompany the carbapenems.

Table 1 also indicates the overall approach to designing longer MDR-TB regimens for adults and children based on the revised grouping. The regimen is designed by adding medicines sequentially going down the three groups.

Apart from the ranking by balance of effectiveness and harms, choice is also determined by: a preference for oral over injectable agents; the results of drug-susceptibility testing (DST); the reliability of existing DST methods; population drug resistance levels; history of previous use of the medicine in a patient; drug tolerability; and potential drug-drug interactions.

Consultation on how best to optimise these aspects of MDR-TB treatment is ongoing. This includes the minimum number of medicines required in designing MDR-TB regimens based on the revised grouping, while maximising regimen efficacy in the presence of resistance to or tolerability of individual agents.

Options for the choice of agents for the intensive and continuation phases, more detailed guidance on patient selection criteria, number of medicines and duration of treatment, adult and paediatric dosing, treatment of extensively drug resistant disease (XDR-TB), and the use of DST results will be provided at the time of release of the final WHO guidelines.

Table 1. Grouping of medicines recommended for use in longer MDR-TB regimens

GROUP	MEDICINE	Abbreviation
Group A: Include all three medicines (unless they cannot be used)	Levofloxacin <u>OR</u> Moxifloxacin Bedaquiline ^{1,4} Linezolid ²	Lfx Mfx Bdq Lzd
Group B: Add both medicines (unless they cannot be used)	Clofazimine Cycloserine <u>OR</u> Terizidone	Cfz Cs Trd
Group C: Add to complete the regimen and when medicines from Groups A and B cannot be used	Ethambutol Delamanid ^{3,4} Pyrazinamide ⁵ Imipenem-cilastatin <u>OR</u> Meropenem ⁶ Amikacin (<u>OR</u> Streptomycin) ⁷ Ethionamide <u>OR</u> Prothionamide <i>p</i> -aminosalicylic acid	E Dlm Z Ipm-Cln Mpm Am (S) Eto Pto PAS

1. Evidence on the safety and effectiveness of Bdq beyond 6 months was insufficient for review; extended Bdq use in individual patients will need to follow '[off-label' use best practices](#)'.
2. Optimal duration of use of Lzd is not established. Use for at least 6 months was shown to be highly effective, although toxicity may limit its use.

3. The position of Dlm will be re-assessed once individual patient data from trial 213 has been reviewed; these data were not available for the evidence assessment in July outlined above. Evidence on the safety and effectiveness of Dlm beyond 6 months was insufficient for review; extended use of Dlm in individual patients will need to follow '[off-label' use best practices](#)'.
4. Evidence on concurrent use of Bdq and Dlm was insufficient for review.
5. Z is only counted as an effective agent when DST results confirm susceptibility.
6. Amoxicillin-Clavulanic acid is administered with every dose of Imp-Cln or Mpm but is not counted as a separate agent and should not be used as a separate agent.
7. Am and S are only to be considered if DST results confirm susceptibility and high-quality audiology monitoring for hearing loss can be ensured. S is to be considered only if Am cannot be used and if DST results confirm susceptibility (S resistance is not detectable with 2nd line molecular line probe assays and phenotypic DST is required).

Shorter MDR-TB regimen^{iv}

- The STREAM Stage 1 trial showed that, in eligible patients, treatment success was similar between patients receiving a shorter MDR-TB regimen and longer regimens conforming to prior WHO recommendations.
- In observational studies, shorter MDR-TB regimens similar to the one studied in the STREAM Stage 1 trial showed an overall comparable likelihood of treatment success with longer regimens, with a lower risk of treatment interruption. However, shorter regimens were associated with higher risk of treatment failure and relapse compared to longer regimens, especially when resistance to key medicines in the shorter regimen was present or when longer regimens included one or more of the Group A medicines listed above in Table 1.
- Evidence is lacking for the performance of shorter MDR-TB regimens modified from the standardized form recommended in 2016 (e.g. bedaquiline or linezolid replacing the injectable agent or levofloxacin replacing moxifloxacin).

Choice of a MDR-TB regimen

- Treatment options for MDR-TB are increasingly becoming more individualised as a result of innovations in diagnostics and growing scientific understanding of the molecular basis for drug resistance and the pharmacokinetics and pharmacodynamics of TB medicines. Three signals are clear from the current scientific evidence assessment:
 - The feasibility of effective and **fully oral treatment regimens** for most patients;

The need to ensure that drug resistance is excluded (at least to the fluoroquinolones and

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_25564

