

EBOLA VIRUS DISEASE

Democratic Republic of the Congo

External Situation Report 08



EBOLA VIRUS DISEASE

Democratic Republic of the Congo

External Situation Report 8

Date of issue:
25 September 2018
Data as reported by: 23 September 2018

1. Situation update



The Ebola virus disease (EVD) outbreak in North Kivu and Ituri provinces of the Democratic Republic of the Congo continues to be closely monitored, with the Ministry of Health (MoH), WHO and partners making progress in response to the outbreak. Recent trends (Figure 1) suggest that control measures are working, although these trends must be interpreted with caution. The outbreak remains ongoing in Beni, Mabalako and Butembo health zones in North Kivu. Tchomia Health Zone in Ituri Province was newly affected in the past week. Additional risks remain high following the movement of several cases in recent weeks across health zones, including Kalunguta.

Since our last situation report on 18 September 2018 (*External Situation Report 7*), an additional eight (8) new confirmed EVD cases and three (3) deaths (1 each in Beni, Butembo and Tchomia) have been reported. The new confirmed cases were reported from four health zones: Beni (4), Mabalako (1), Tchomia (2) and Butembo (1). This is the first time confirmed cases have been reported from Tchomia Health Zone in Ituri which borders Uganda. Both confirmed cases reported from Tchomia were linked to the ongoing Beni transmission chain.

As of 23 September 2018, 9 suspected cases have been identified following the investigation of 17 alerts. Table 1 summarizes the geographical distribution of these suspected cases. No additional cases were reported among health workers in the last week. Cumulatively, 19 health workers have been affected (18 confirmed and one probable), three of whom have died. All health workers' exposures occurred in health facilities outside dedicated Ebola treatment centres (ETCs).

As of 23 September 2018, a total of 150 confirmed and probable EVD cases, including 100 deaths, have been reported resulting in a case fatality ratio (CFR) of 67%. Of the 150 cases, 119 are confirmed and 31 are probable. Among confirmed cases only, the CFR was 58% (69/119). Of the 140 cases with known age and sex, 56% (n=78) are female. Among females, the most affected age group is 25-34 years, while among men, the most affected age group is 35-44 years (Figure 2).

As of 23 September 2018, 41 confirmed cases have recovered and been discharged from ETCs; eight (8) confirmed cases remain hospitalized in ETCs.

The epicentres of the outbreak remain in Mabalako and Beni health zones in North Kivu Province, reporting 60% (n=90) and 22% (n=33) of all confirmed and probable cases, respectively. However, since late August 2018, most new cases have occurred in Beni or are related to the ongoing Beni transmission chain. Of the total deaths reported to date, 65% (n=65) were from Mabalako, while 24% (n=24) were from Beni (Table 1 and Figure 3). Additionally, five other health zones in North Kivu Province and three in Ituri Province have reported confirmed and probable cases (Table 1 and Figure 3).

The MoH, WHO and partners are monitoring and investigating all alerts in affected areas, in other provinces in the Democratic Republic of the Congo (including Kisangani and Tshopo provinces) and in neighbouring countries. As of 23 September 2018, nine (9) suspected cases are awaiting laboratory testing within outbreak


affected areas (Table 1). Since the last report was published, alerts were investigated in several provinces of the Democratic Republic of the Congo as well as in Uganda and South Sudan; and to date, EVD has been ruled out in all alerts from neighbouring provinces and countries.

Table 1: Ebola virus disease cases by classification and health zones in North Kivu and Ituri provinces, Democratic Republic of the Congo, as of 23 September 2018

Case classification/ status	North Kivu							Ituri			Total
	Beni	Butembo	Oicha	Mabalako	Musienene	Masereka	Kalunguta	Komanda	Mandima	Tchomia	
Probable*	4	2	1	21	1	0	0	0	2	0	31
Confirmed	29	6	2	69	0	1	1	0	9	2	119
Total confirmed and probable	33	8	3	90	1	1	1	0	11	2	150
Suspected cases currently under investigation	3	1	1	4	0	0	0	0	0	0	9
Deaths											
Total deaths	24	4	1	65	1	1	0	0	3	1	100
Deaths in confirmed cases	20	2	0	44	0	1	0	0	1	1	69


*Includes n=27 community deaths, retrospectively identified from clinical records, tentatively classified as probable cases pending further investigation.

Figure 1. Confirmed and probable Ebola virus disease cases by week of illness onset, data as of 23 September 2018 (n=150)*




*Case counts in recent weeks may be incomplete due to reporting details. All trends should be interpreted with caution.

Figure 2: Confirmed and probable Ebola virus disease cases by age and sex, North Kivu and Ituri provinces, Democratic Republic of the Congo, 23 September 2018 (n=140)



*Age/sex is currently unknown for n=10 cases.

Figure 3: Geographical distribution of confirmed and probable Ebola virus disease cases in North Kivu and Ituri provinces, Democratic Republic of the Congo, 23 September 2018 (n=150)



Context

North Kivu and Ituri are among the most populated provinces in the Democratic Republic of the Congo. North Kivu shares borders with Uganda and Rwanda. The provinces have been experiencing intense insecurity and a worsening humanitarian crisis, with over one million internally displaced people and a continuous movement of refugees to neighbouring countries, including Uganda, Burundi and Tanzania. The Democratic Republic of the Congo is also experiencing multiple disease outbreaks, including three separate outbreaks of circulating vaccine-derived poliovirus type 2 (cVDPV2) in the provinces of Ituri, Mongola, Maniema and Haut Lomami, Tanganyika and Haut Katanga, and outbreaks of cholera, measles and monkeypox spread across the country.

Current risk assessment

This outbreak of EVD is affecting north-eastern provinces of the Democratic Republic of the Congo, which border Uganda, Rwanda and South Sudan. Potential risk factors for transmission of EVD at the national and regional levels include the transportation links between the affected areas, the rest of the country, and neighbouring countries; the internal displacement of populations; and the displacement of Congolese refugees to neighbouring countries. The country is concurrently experiencing other epidemics (e.g. cholera, vaccine-derived poliomyelitis), and a long-term humanitarian crisis. Additionally, the security situation in North Kivu and Ituri may hinder the implementation of response activities. Based on this context, the public health risk was assessed to be high at the national and regional levels, and low globally.

As the risk of national and regional spread remains high, it is important for neighbouring provinces and countries to continue to enhance surveillance and preparedness activities. WHO will continue to work with neighbouring countries and partners to ensure health authorities are alerted and are operationally ready to respond

Strategic approach to the prevention, detection and control of EVD

WHO recommends the implementation of proven strategies for the prevention and control EVD outbreaks. These include (i) strengthening the multi-sectoral coordination of the response, (ii) enhanced surveillance, including active case finding, case investigation, contact tracing and surveillance at Points of Entry (PoE), (iii) strengthening diagnostic capabilities, (iv) case management, (v) infection prevention and control in health facilities and communities, including safe and dignified burials, (vi) risk communication, social mobilization and community engagement, (vii) psychosocial care (viii) vaccination of risk groups (ix) research and (x) operational support and logistics.

2. Actions to date

Coordination of the response

- ⇒ Field activities were suspended in Beni this Sunday 23 September 2018 following clashes between rebels and the Congolese armed forces which took place on 22 September 2018. Response activities will resume as soon as the calm has returned to the city.
- ⇒ This attack underscores the challenges faced by the government and health organisations when responding to an Ebola outbreak in a conflict zone.

Surveillance

- As of 23 September 2018, a total of 1836 contacts remain under surveillance. Overall, there was an improvement in the proportion of contacts followed up daily, with 93-98% seen and monitored over the past week compared to 85-94% during the preceding week. This improvement should be interpreted carefully considering that 5 of the confirmed cases reported over the past week were not registered as contacts. However, as field activities were suspended in Beni on 23 September, the proportion of contacts followed up in Beni fell to 35% on that day. This highlights a need to reinforce contact listing and tracing of missing contacts who could eventually generate new transmission chains.
- As of 22 September, more than 5 million travellers have been screened at PoEs, and over 17,000 means of travel have been decontaminated.

Laboratory

- MSF will provide a Piccolo blood chemistry analyser in Butembo to support biochemistry analyses.
- Laboratory testing capacity for Ebola has been established in hospital facilities in Beni, Goma, Mangina and Butembo to facilitate rapid diagnosis of suspected cases.

Case management

- There are currently four ETCs in the country. The ETC supported by International Medical Corps (IMC) is in Makeke Health Area (Ituri Province), two additional ETCs are operational in Beni and Mangina with support from The Alliance for International Medical Action (ALIMA) and Médecins Sans Frontières (MSF), respectively and MSF Switzerland is collaborating with the MoH to support the ETC in Butembo. Assessments for potential additional locations for ETCs are ongoing.
- Development of a proposal for a 1-year Ebola survivor programme is underway, including clinical management of complications associated with Ebola illness, screening and prevention of sexual secondary transmission, and psycho-social support.
- ETCs continue to provide therapeutics under the monitored emergency use of unregistered and experimental interventions (MEURI) protocol in collaboration with the MoH and the Institut National de Recherche Biomédicale (INRB). WHO is providing technical clinical expertise onsite and is assisting with the creation of a data safety management board.
- As of 23 September 2018, 38 patients have received investigational Ebola therapeutics in addition to optimized standard of care, following informed consent within the context of compassionate use protocols. Of these patients, 19 have been discharged, 7 are still under treatment and 12 have died. The median delay between onset of symptoms and beginning treatment is 5 days.

- ⇒ Trainings of health workers in infection prevention and control (IPC) practices including both standard precautions as well as recognition of signs and symptoms of Ebola with immediate triage are ongoing in affected areas including Beni, Butembo, Kisangani, Komanda, Butsili, Mabolio, Mangina and Tchomia.
- ⇒ Routine water, sanitation and hygiene (WASH) activities are ongoing.
- ⇒ In addition to many other structures that have been previously decontaminated, additional health structures that previously admitted confirmed EVD cases were decontaminated in Tchomia, Mangina and Mandradelé.
- ⇒ Following the confirmed EVD positive case in Tchomia, an evaluation of WASH and IPC needs has been conducted at the General Reference Hospital of Tchomia. IPC activities are continuing in this area.
- ⇒ Safe and dignified burials (SDBs) continue to be conducted by Red Cross SDB teams in Mabalako, Beni and Butembo. Due to the confirmed EVD case in Ituri, the Red Cross is strengthening the SDB team in Bunia. In addition, SDB team training is being conducted in Goma to increase preparedness.
- ⇒ As part of the Red Zone Strategy, the Red Cross together, with MoH and WHO, has trained Government Civil protection teams from Beni, Butembo and Oicha to be able to respond in Red Zone areas.

Implementation of ring vaccination protocol

- ⇒ As of 22 September, 11,417 people consented and were vaccinated against Ebola: 4,056 in Mabalako, 3,652 in Beni, 1,632 in Mandima, 834 in Butembo, 613 in Katwa, 270 in Masereka, 164 in Komanda, 121 in Oicha, 65 in Kinshasa (health workers to be deployed) and 10 in Tchomia.
- ⇒ As of 22 September, 8 vaccination rings were active: 3 in Beni (including two new rings), 2 in Butembo (including 1 new ring), 1 in Katwa, 1 in Tchomia (new ring) and 1 in Mabalako (new ring).
- ⇒ As of 25 September, 63 vaccination rings have been defined, including 26 rings of health and other frontline workers. These rings notably include the contacts (and their contacts) of the confirmed cases from the last four weeks.

Psychosocial care

- ⇒ Routine psychosocial activities are provided in ETCs, to affected people and families, to contacts and to orphans. Coverage for each targeted population is 100%, except for among contacts (57%).

Risk communication, social mobilization and community engagement

- The MoH, WHO, UNICEF, the Red Cross and partners are intensifying activities to engage with local community leaders and community networks in the affected areas. While public awareness activities are ongoing through community radio and social mobilization in public places, such as markets, the priority is to engage community groups in the Ebola response.
- In Beni and Butembo, teams are working to address community incidents caused as a result of reluctance or refusal to comply with Ebola response teams. In Butembo, community engagement was strengthened through collaboration with a popular singer, Mayaya Santa, to produce a song with key messages.

Logistics

- An operations hub has been established in Butembo with dedicated coordination support from WHO, partners, and the MoH.

Resource mobilization

- Implementation of and resource mobilization for the joint strategic response plan, approved by the Minister of Health of the Democratic Republic of the Congo, is progressing well, in collaboration with the national authorities and all partners.

Preparedness

- The WHO Regional Office for Africa has updated the regional preparedness plan and reprioritized neighbouring countries based on proximity to North Kivu, the current EVD epicentre. The new prioritizations are as follows: Priority 1: Rwanda, Uganda, South Sudan and Burundi; Priority 2: Angola, Congo, Central African Republic, Tanzania, Zambia. These countries were prioritised based on their proximity to the EVD epicentre and their capacity to respond to the disease.

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_25598

