

WHO TobLabNet
Official Method
SOP 07

STANDARD OPERATING PROCEDURE FOR DETERMINATION OF AMMONIA IN CIGARETTE TOBACCO FILLER

Tobacco Free Initiative
Tobacco Laboratory Network (TobLabNet)



**World Health
Organization**



**WHO TobLabNet
Official Method
SOP 07**

**Standard operating procedure
for determination of ammonia
in cigarette tobacco filler**



WHO Library Cataloguing-in-Publication Data

Standard operating procedure for determination of ammonia in cigarette tobacco filler, WHO TobLabNet Official Method SOP07.

1.Tobacco Products – analysis. 2.Smoke – analysis. 3.Excipients – toxicity. 4.Ammonia - toxicity. 5.Toxicity Tests. I.World Health Organization. Tobacco Laboratory Network.

ISBN 978 92 4 151605 1

(NLM classification: QV 137)

This publication was originally published under ISBN 978 92 4 151046 2

© World Health Organization 2016

All rights reserved. Publications of the World Health Organization are available on the WHO website (<http://www.who.int>) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; email: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications – whether for sale or for non-commercial distribution – should be addressed to WHO Press through the WHO website (http://www.who.int/about/licensing/copyright_form/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Printed in Switzerland



No.: SOP 07
Date: July 2016



**World Health
Organization**

**World Health Organization
Tobacco Laboratory Network**

Standard operating procedure for method

**Determination of ammonia
in cigarette tobacco filler**

Method:	Determination of ammonia in cigarette tobacco filler
Analytes:	Ammonia (CAS # 7664-41-7)
Matrix:	Cigarette tobacco filler
Last update:	July 2016



No.: SOP 07

Date: July 2016

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

No machine smoking regimen can represent all human smoking behaviour: machine smoking testing is useful for characterizing cigarette emissions for design and regulatory purposes, but communication of machine measurements to smokers can result in misunderstanding about differences between brands in exposure and risk. Data on smoke emissions from machine measurements may be used as inputs for product hazard assessment, but they are not intended to be nor are they valid as measures of human exposure or risk. Representing differences in machine measurements as differences in exposure or risk is a misuse of the results of testing with WHO TobLabNet-recommended methods.



No.: SOP 07

Date: July 2016

FOREWORD

This document was prepared by members of the World Health Organization (WHO) Tobacco Laboratory Network (TobLabNet) as an analytical method standard operating procedure (SOP) for measuring ammonia in cigarette tobacco filler.

INTRODUCTION

In order to establish comparable measurements for testing tobacco products globally, consensus methods are required for measuring specific contents and emissions of cigarettes. The Conference of the Parties to the WHO Framework Convention on Tobacco Control (WHO FCTC) at its third session, in Durban, South Africa, in November 2008, “recalling its decisions FCTC/COP1(15) and FCTC/COP2(14) on the elaboration of guidelines for implementation of Articles 9 (Regulation of the contents of tobacco products) and 10 (Regulation of tobacco product disclosures) of the WHO FCTC, noting the information contained in the report of the working group to the third session of the Conference of the Parties on the progress of its work ... requested the Convention Secretariat to invite WHO’s Tobacco Free Initiative to ... validate, within five years, the analytical chemical methods for testing and measuring cigarette contents and emissions” (FCTC/COP/3/REC/1).

Using the criteria for prioritization set at its third meeting in Ottawa, Canada, in October 2006, the working group on Articles 9 and 10 identified the following contents for which methods for testing and measuring (analytical chemistry) should be validated as a priority:

- nicotine
- ammonia
- propylene glycol (propane-1,2-diol)
- glycerol (propane-1,2,3-triol)
- triethylene glycol (2,2-ethylenedioxybis(ethanol)).

Measurement of these contents will require validation of three methods: one for nicotine, one for ammonia and one for humectants.

Using the criteria for prioritization set at the meeting in Ottawa mentioned above, the working group identified the following emissions in mainstream smoke for which methods for testing and measurement (analytical chemistry) should be validated as a priority:

- 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)
- N-nitrosornicotine (NNN)



- acetaldehyde
- acrylaldehyde (acrolein)
- benzene
- benzo[a]pyrene
- 1,3-butadiene
- carbon monoxide
- formaldehyde.

Measurement of these emissions with the two smoking regimens described below will require validation of five methods: one for tobacco-specific nitrosamines (NNK and NNN), one for benzo[a]pyrene, one for aldehydes (acetaldehyde, acrolein and formaldehyde), one for volatile organic compounds (benzene and 1,3-butadiene) and one for carbon monoxide.

The table below sets out the two smoking regimens for validation of the test methods referred to above.

Smoking regimen	Puff volume (mL)	Puff frequency	Filter ventilation holes
ISO regimen: ISO 3308: <i>Routine analytical cigarette smoking machine – definitions and standard conditions</i>	35	Once every 60 s	No modification
Intense regimen: Same as ISO 3308, but modified as indicated	55	Once every 30 s	All ventilation holes must be blocked 100% as described in WHO TobLabNet SOP 01.

This SOP was prepared to describe the procedure for the determination of ammonia in cigarette tobacco filler

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_26844

