


Integrated Management of Pregnancy and Childbirth

Pregnancy, Childbirth, Postpartum and Newborn Care: A guide for essential practice



Integrated Management of Pregnancy and Childbirth

Pregnancy, Childbirth, Postpartum and Newborn Care: A guide for essential practice

Third Edition



**World Health
Organization**

WHO Library Cataloguing-in-Publication Data

Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice – 3rd ed.

1.Labor, Obstetric. 2.Delivery, Obstetric. 3.Prenatal Care. 4.Perinatal Care. 5.Postnatal Care. 6.Pregnancy Complications. 7.Practice Guideline. I.World Health Organization. II.UNFPA. III.UNICEF. IV.World Bank.

ISBN 978 92 4 154935 6

(NLM classification: WQ 176)

© **World Health Organization 2015**

All rights reserved. Publications of the World Health Organization are available on the WHO website (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications –whether for sale or for non-commercial distribution– should be addressed to WHO Press through the WHO website (www.who.int/about/licensing/copyright_form/en/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Printed in Luxembourg.

PREFACE

'Pregnancy, Childbirth, Postpartum and Newborn Care: A guide for essential practice' (PCPNC) has been updated to include recommendations from recently approved WHO guidelines relevant to maternal and perinatal health. These include pre-eclampsia & eclampsia; postpartum haemorrhage; postnatal care for the mother and baby; newborn resuscitation; prevention of mother-to-child transmission of HIV; HIV and infant feeding; malaria in pregnancy, tobacco use and second-hand exposure in pregnancy, post-partum depression, post-partum family planning and post abortion care.

This revised guide brings a full range of updated evidence – based norms and standards that enable health care providers at the first health care level to provide high-quality, integrated care during pregnancy and childbirth and after birth, both for mothers and babies. This guide will support countries in their efforts to reach every woman and child and ensure that pregnancy, birth and the first postnatal weeks are the joyful and safe experience they should be. The guide will be updated periodically as new WHO recommendations become available.

This guide represents a common understanding between WHO, UNFPA, UNICEF, and the World Bank of key elements of an approach to reducing maternal and perinatal mortality and morbidity. These agencies co-operate closely in efforts to reduce maternal and perinatal mortality and morbidity. The principles and policies of each agency are governed by the relevant decisions of each agency's governing body and each agency implements the interventions described in this document in accordance with these principles and policies and within the scope of its mandate.

Acknowledgements

ACKNOWLEDGEMENTS

The 2006 edition was prepared by a team of the World Health Organization, Department of Reproductive Health and Research (RHR), led by Jerker Liljestrand and Jelka Zupan. The concept and first drafts were developed by Sandra Gove and Patricia Whitesell/ACT International, Atlanta, Jerker Liljestrand, Denise Roth, Betty Sweet, Anne Thompson, and Jelka Zupan. Revisions were subsequently carried out by Annie Portela, Luc de Bernis, Ornella Lincetto, Rita Kabra, Maggie Usher, Agostino Borra, Rick Guidotti, Elisabeth Hoff, Mathews Matthai, Monir Islam, Felicity Savage, Adepeyu Olukoya, Aafje Rietveld, TinTin Sint, Ekpini Ehounou, Suman Mehta.

Valuable inputs were provided by WHO Regional Offices and WHO departments:

- Reproductive Health and Research (RHR)
- Maternal, Newborn, Child and Adolescent Health (MCA)
- HIV/AIDS
- Nutrition for Health and Development (NHD)
- Essential Medicines and Health Products (EMP)
- Immunization, Vaccines and Biologicals (IVB)
- Mental Health and Substance Abuse (MSD)
- Gender, Women and Health (GWH)
- Prevention of Blindness and Deafness (PBD)
- Prevention of Noncommunicable Diseases (PND)

Editing: Nina Mattock, Richard Casna

Layout: rsdesigns.com sàrl

Cover design: Máire Ní Mhearáin

WHO acknowledges with gratitude the generous contribution of over 100 individuals and organizations in the field of maternal and newborn health, who took time to review the 2006 version at different stages of its development. They came from over 35 countries and brought their expertise and wide experience to the final text.

The guide was also reviewed and endorsed by the International Confederation of Midwives, the International Federation of Gynecology and Obstetrics and International Pediatric Association.



International
Confederation of Midwives



International Federation of
Gynecology and Obstetrics



International
Pediatric Association

The financial support towards the preparation and production of the 2006 version was provided by UNFPA, the World Bank and the Governments of Australia, Japan and the United States of America.

The present version was revised and updated by Jelka Zupan and Juana Willumsen. Overall technical direction and supervision of the document was maintained by Maurice Bucagu of WHO/MCA.

Technical inputs were provided by Matthews Mathai, Annie portela, severin von Xylander, Metin Ahmet Gulmezoglu, Bela Ganatra, Mercedes Semenas Bonet, Taiwo Olufemi Oladapo, Ehizibue Peter Olumese, Susheela Engelbrecht, Fabio Uxa, Patricia Coffey, Satvinder Singh, Mustafa Shaffiq Essajee, Ahmadu Yakubu. Financial support for the update was provided by The Norwegian Agency for Development Cooperation and Ministry of Foreign Affairs and International Development (France).

TABLE OF CONTENTS

INTRODUCTION

- i Introduction
- i2 How to read the Guide
- i3 Structure and presentation
- i4 Assumptions underlying the guide

A PRINCIPLES OF GOOD CARE

- A2 Communication
- A3 Workplace and administrative procedures
- A4 Standard precautions and cleanliness
- A5 Organising a visit

B QUICK CHECK, RAPID ASSESSMENT AND MANAGEMENT OF WOMEN OF CHILDBEARING AGE

- B2 Quick check
- B3-B7 Rapid assessment and management
 - B3 Airway and breathing
 - B3 Circulation (shock)
 - B4-B5 Vaginal bleeding
 - B6 Convulsions or unconscious
 - B6 Severe abdominal pain
 - B6 Dangerous fever
 - B7 Labour
 - B7 Other danger signs or symptoms
 - B7 If no emergency or priority signs, non urgent

B EMERGENCY TREATMENTS FOR THE WOMAN

- B9 Airway, breathing and circulation
 - B9 Manage the airway and breathing
 - B9 Insert IV line and give fluids
 - B9 If intravenous access not possible
- B10-B12 Bleeding
 - B10 Massage uterus and expel clots
 - B10 Apply bimanual uterine compression
 - B10 Apply aortic compression
 - B10 Give oxytocin
 - B10 Give misoprostol
 - B10 Give ergometrine
 - B11 Remove placenta and fragments manually
 - B11 After manual removal of placenta
 - B12 Repair the tear or episiotomy
 - B12 Empty bladder
- B13-B14 Important considerations in caring for a woman with eclampsia or pre-eclampsia
 - B13 Give magnesium sulphate
 - B13 Important considerations in caring for a woman with eclampsia
 - B14 Give diazepam
 - B14 Give appropriate antihypertensive drug
- B15 Infection
 - B15 Give appropriate IV/IM antibiotics
- B16 Malaria
 - B16 Treatment of uncomplicated P.falciparum malaria in pregnancy
- B17 Refer the woman urgently to the hospital
 - B17 Essential emergency drugs and supplies for transport and home delivery

B BLEEDING IN EARLY PREGNANCY AND POST-ABORTION CARE

- B19 Examination of the woman with bleeding in early pregnancy and post-abortion care
- B20 Give preventive measures
- B21 Advise and counsel on post-abortion care
 - B21 Advise on self-care
 - B21 Advise and counsel on family planning
 - B21 Provide information and support after abortion
 - B21 Advise and counsel during follow-up visits

Table of contents

C ANTENATAL CARE

- C2** Assess the pregnant woman: pregnancy status, birth and emergency plan
 - C3** Check for pre-eclampsia
 - C4** Check for anaemia
 - C5** Check for syphilis
 - C6** Check for HIV status
- C7** Respond to observed signs or volunteered problems
 - C7** If no fetal movement
 - C7** If ruptured membranes and no labour
 - C8** If fever or burning on urination
 - C9** If vaginal discharge
 - C10** If signs suggesting HIV infection
 - C10** If smoking, alcohol or drug abuse, or history of violence
 - C11** If cough or breathing difficulty
 - C11** If taking antituberculosis drugs
- C12** Give preventive measures
- C13** Advise and counsel on nutrition and self-care
- C14-C15** Develop a birth and emergency plan
 - C14** Facility delivery
 - C14** Home delivery with a skilled attendant
 - C15** Advise on labour signs
 - C15** Advise on danger signs
 - C15** Discuss how to prepare for an emergency in pregnancy
- C16** Advise and counsel on family planning
 - C16** Counsel on the importance of family planning
 - C16** Special consideration for family planning counselling during pregnancy
- C17** Advise on routine and follow-up visits
- C18** Home delivery without a skilled attendant

D CHILDBIRTH - LABOUR, DELIVERY AND IMMEDIATE POSTPARTUM CARE

- D2** Examine the woman in labour or with ruptured membranes
- D3** Decide stage of labour
- D4-D5** Respond to obstetrical problems on admission
- D6-D7** Give supportive care throughout labour
 - D6** Communication
 - D6** Cleanliness
 - D6** Mobility
 - D6** Urination
 - D6** Eating, drinking
 - D6** Breathing technique
 - D6** Pain and discomfort relief
 - D7** Birth companion
- D8-D9** First stage of labour
 - D8** Not in active labour
 - D9** In active labour
- D10-D11** Second stage of labour: deliver the baby and give immediate newborn care
- D12-D13** Third stage of labour: deliver the placenta
- D14-D18** Respond to problems during labour and delivery
 - D14** If fetal heart rate <120 or >160 beats per minute
 - D15** If prolapsed cord
 - D16** If breech presentation
 - D17** If stuck shoulders (Shoulder dystocia)
 - D18** If multiple births
- D19** Care of the mother and newborn within first hour of delivery of placenta
- D20** Care of the mother one hour after delivery of placenta
- D21** Assess the mother after delivery
- D22-D25** Respond to problems immediately postpartum
 - D22** If vaginal bleeding
 - D22** If fever (temperature >38°C)
 - D22** If perineal tear or episiotomy (done for lifesaving circumstances)
 - D23** If elevated diastolic blood pressure
 - D24** If pallor on screening, check for anaemia
 - D24** If mother severely ill or separated from the child
 - D24** If baby stillborn or dead
- D25** Give preventive measures

https://www.yunbaogao.cn/report/index/report?reportId=5_27566

预览已结束，完整报告链接和二维码如下：



POSTPARTUM CARE (CONTINUED)

Postpartum

(s)
ms

F PREVENTIVE MEASURES AND ADDITIONAL TREATMENTS FOR THE WOMAN

F2-F4 Preventive measures

- F2 Give tetanus toxoid
- F2 Give iron and folic acid
- F2 Give mebendazole
- F3 Give aspirin and calcium (if in area of low dietary calcium intake)
- F3 Motivate on adherence with treatments
- F4 Give preventive intermittent treatment for falciparum malaria in pregnancy
- F4 Advise to use insecticide-treated bednet
- F4 Give appropriate oral antimalarial treatment (uncomplicated P. falciparum malaria)
- F4 Give paracetamol

F5-F6 Additional treatments for the woman

- F5 Give appropriate oral antibiotics
- F6 Give benzathine penicillin IM
- F6 Observe for signs of allergy