

Report on global sexually transmitted infection surveillance 2013



**Report on global sexually transmitted
infection surveillance
2013**

WHO Library Cataloguing-in-Publication Data :

Report on global sexually transmitted infection surveillance 2013.

1. Sexually Transmitted Diseases – epidemiology. 2. Epidemiological Monitoring. 3. Epidemiologic Methods. I. World Health Organization.

ISBN 978 92 4 150740 0

(NLM classification: WC 140)

© World Health Organization 2014

All rights reserved. Publications of the World Health Organization are available on the WHO website (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications –whether for sale or for non-commercial distribution– should be addressed to WHO Press through the WHO website (www.who.int/about/licensing/copyright_form/en/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Printed by the WHO Document Production Services, Geneva, Switzerland

Acknowledgments

The World Health Organization (WHO) would like to express gratitude to the Member States working to collect and share data on the burden of sexually transmitted infections (STIs) in their countries. It is the work of dedicated individuals at the local and national levels that is the foundation for this report.

WHO's Department of Reproductive Health and Research would like to acknowledge the WHO regional staff members who support HIV and STI surveillance and programmes, including: Monica Alonso, Emil Asamoah-Odei, Hamida Khattabi, Lali Khotenashvili, Ying-Ru Lo, Assimawe Pana, Razia Pendse, Gabriele Riedner, Suzanne Serruya, Annemarie Stengaard and Dongbao Yu. In addition, we would like to thank our colleagues in the Departments of HIV/AIDS and Reproductive Health and Research for their support in producing this baseline report and in the broader efforts to strengthen STI surveillance: Maria Babovic, Michel Beusenberg, Txema Garcia Calleja, Chika Hayashi, Gottfried Hirnschall, Irene Martin and Gundo Weiler.

We also would like to express our gratitude to the WHO Gonococcal Antimicrobial Surveillance Programme (GASP) regional collaborating centres and technical leads: Manju Bala, Jo Anne Dillon, Amina Hancali, Catherine Ison, Sarah Kidd, Monica Lahra, David Lewis, Pilar Ramon Pardo, Magnus Unemo and Tom Wong.

We would like to thank Gianfranco Spiteri and the European Centre for Disease Prevention and Control (ECDC) for their collaboration in efforts to strengthen STI surveillance in Europe.

We thank our consultant, Richard Steen, who prepared much of the analysis and drafted this report. In addition, Iyanathi Abeyewikreme, Angel Manuel Alvarez, Angkana Charoenwatanachokchai, Xiang-Sheng Chen, Maura Laverty, Yang Ligang, Francis Ndowa and Saman Wijesooriya contributed to the case studies shared in this report.

We also wish to thank, at the United States Centers for Disease Control and Prevention (CDC), Gail Bolan, Mary Kamb and Hillard Weinstock for review and input into STI surveillance strengthening and for ensuring that STI surveillance remains a priority. We are grateful also for financial support from the CDC and for the technical contributions of other CDC staff members dedicated to strengthening STI surveillance at the country, regional and global levels.

Lori Newman and Teodora Wi in the WHO Department of Reproductive Health and Research coordinated development of this document. Nathalie Broutet, Marleen Temmerman and Igor Toskin contributed further input and review.

We thank Green Ink (United Kingdom) for editing.

Contents

Executive summary	1
Introduction	4
1. Progress controlling mother-to-child transmission of syphilis	6
1.1 Syphilis in pregnancy	6
Indicator 1: Percentage of ANC attendees who were tested for syphilis	8
Indicator 2: Percentage of ANC attendees tested who were positive for syphilis	8
Indicator 3: Percentage of ANC attendees positive for syphilis who received treatment	9
Box 1: Using routinely reported data to monitor progress towards EMTCT of syphilis	9
1.2 Congenital syphilis	10
Indicator 4: Number of reported congenital syphilis cases	11
Box 2: Elimination of congenital syphilis in the WHO European Region	11
Box 3: A tool to estimate the burden of syphilis in pregnancy and adverse outcomes	12
2. Slowing sexual transmission – infectious syphilis and other STIs	13
2.1 Infectious syphilis	13
Indicator 5: Number of adults reported with syphilis	13
Box 4: Syphilis trends in the WHO Western Pacific Region	15
Box 5: Disaggregating syphilis cases in Mongolia	16
Box 6: Syphilis elimination in China	16
2.2 Genital ulcer disease	18
Indicator 6: Number of adults reported with genital ulcer disease	18
Box 7: Identifying areas of increasing sexual transmission in Cambodia	18
2.3 Urethral discharge and gonorrhoea in men	20
Indicator 7: Number of men reported with urethral discharge	20
Indicator 8: Number of men reported with gonorrhoea	21
3. STI control with key populations	22
3.1 STI control with female sex workers	22
Indicator 9: Percentage of female sex workers (FSWs) with active syphilis	23
Box 8: Evidence of declining syphilis prevalence among sex workers in Asia	24
Box 9: Low syphilis prevalence among sex workers in the Philippines	24
3.2 STI control for men who have sex with men and transgender persons	25
Indicator 10: Prevalence of syphilis among men who have sex with men	25
Box 10: Indonesia: monitoring syphilis among key populations	26

4. Building reliable reporting systems	27
4.1 Assessing and strengthening routine STI reporting	27
Box 11: Efforts in the Republic of Moldova and the WHO European Region to strengthen antenatal STI screening and reporting	28
4.2 Using syndromic information together with etiologic data to improve programmes	28
4.3 Piloting new STI indicators for HIV programmes	29
Box 12: STI surveillance in Latin America and the Caribbean – an example of regional progress	30
5. Gonococcal Antimicrobial Surveillance Programme (GASP)	32
5.1 Aims, structure, coordination and participation	32
5.2 Antimicrobial resistance patterns	33
Box 13: STI surveillance and GASP in countries of the WHO African Region	34
Conclusions	35
Annexes	38
Annex 1: Cascade of indicators for elimination of mother-to-child transmission of syphilis using most recent data reported through the Global AIDS Response Progress Reporting (GARPR) system, 2008–2012	39
Annex 2: Syphilis prevalence reported for sex workers and men who have sex with men (MSM) using most recent data reported through the Global AIDS Response Progress Reporting (GARPR) system, 2008–2012	44
Annex 3: Pilot data on new STI indicators from two regions reported through the Global AIDS Response Progress Reporting (GARPR) system, 2012	46
Annex 4: Data collected through WHO Gonococcal Antimicrobial Surveillance Programme (GASP), 2010–2011	48
Annex 5: Data sources, analysis and interpretation	51
References	53

Abbreviations and acronyms

AMR	antimicrobial resistance
ANC	antenatal care
CAREC	Caribbean Epidemiology Center
CDC	United States Centers for Disease Control and Prevention
CISID	Centralized Information System for Infectious Diseases
DTECS	Dual Testing for the Elimination of Congenital Syphilis
ECDC	European Centre for Disease Prevention and Control
EMTCT	elimination of mother-to-child transmission
ESC	extended-spectrum cephalosporin
FSW	female sex worker
GARPR	Global AIDS Response Progress Reporting
GASP	Gonococcal Antimicrobial Surveillance Programme
GUD	genital ulcer disease
HSV	herpes simplex virus
LGV	lymphogranuloma venereum
MSM	men who have sex with men
MTCT	mother-to-child transmission
PAHO	Pan American Health Organization
PID	pelvic inflammatory disease
PMTCT	prevention of mother-to-child transmission
PPT	periodic presumptive treatment
PWID	people who inject drugs
STI	sexually transmitted infection
UD	urethral discharge
UNAIDS	Joint United Nations Programme on HIV/AIDS
UNICEF	United Nations Children's Fund
WHO	World Health Organization

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_27825

