



Maternal Mortality in 2000:

Estimates developed by
WHO, UNICEF, UNFPA

unicef 



World Health Organization, Geneva

Maternal Mortality in 2000:

Estimates developed by
WHO, UNICEF and UNFPA



Department of Reproductive Health and Research
World Health Organization, Geneva 2004

WHO Library Cataloguing-in-Publication Data

Maternal mortality in 2000 : estimates developed by WHO, UNICEF and UNFPA

1.Maternal mortality 2.Data collection - methods 3.Models, Statistical I.World Health Organization II.UNICEF III.United Nations Population Fund

ISBN 92 4 156270 6

(NLM classification: WQ 16)

© World Health Organization 2004

All rights reserved. Publications of the World Health Organization can be obtained from Marketing and Dissemination, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel: +41 22 791 2476; fax: +41 22 791 4857; email: bookorders@who.int). Requests for permission to reproduce or translate WHO publications—whether for sale or for noncommercial distribution—should be addressed to Publications, at the above address (fax: +41 22 791 4806; email: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use.

Printed in



Acknowledgements

This document was prepared by Carla AbouZahr^a of WHO and Tessa Wardlaw^b of UNICEF on the basis of a technical paper originally developed by Kenneth Hill and Yoonjoung Choi, Johns Hopkins University. Valuable inputs and assistance were provided by Colin Mathers, Kenji Shibuya, Nyein Nyein Lwin, Ana Betran and Elisabeth Aahman. Particular thanks to Gareth Jones, Paul Van Look and France Donnay for their guidance, advice and unfailing support.

^a Technical officer, Measurement and Health Information Systems, Evidence and Information for Policy, WHO, Geneva.

Correspondence to C. AbouZahr, Evidence and Information for Policy, World Health Organization, Avenue Appia, 1211 Geneva 27, Switzerland.

^b Senior Programme Officer, Statistics and Monitoring, Division of Policy and Planning, UNICEF, New York.

Abbreviations

CEE/CIS	Central and Eastern Europe/Commonwealth of Independent States
DHS	Demographic and Health Surveys
EIP	WHO Evidence and Information for Health Policy Cluster
GFR	general fertility rate
LASSAME	Countries in Latin America and the Caribbean, sub-Saharan Africa and the Middle East
MDG	Millennium Development Goal
MMR	maternal mortality ratio
OECD	Organisation for Economic Co-operation and Development
PMDF	proportion maternal among deaths of women of reproductive age
RAMOS	reproductive age mortality study
TFR	total fertility rate
UN	United Nations
UNFPA	United Nations Population Fund
UNICEF	United Nations Children's Fund
WHO	World Health Organization



Contents

Executive summary	1
Introduction	3
Background	3
Maternal mortality: The measurement challenge	3
Definitions and measures of maternal mortality	3
Why maternal mortality is difficult to measure	4
Approaches for measuring maternal mortality	5
The development of 2000 estimates of maternal mortality	7
Process for developing the 2000 estimates	7
Sources of country data used for the 2000 estimates	7
Differences between the 2000 methodology compared with 1995	10
Analysis and interpretation of 2000 estimates	10
Maternal mortality estimates for 2000	10
Differences between 2000 estimates and nationally reported data	11
Comparing 2000 estimates with those for 1990 and 1995	12
Using the 2000 maternal mortality estimates	13
What can the 2000 estimates be used for?	13
What should they NOT be used for?	13
Why can the 2000 estimates NOT be used to estimate trends?	13
Next steps	14
Generate better information	15

Tables

Table 1	
Sources of maternal mortality data used in developing the 2000 estimates	8
Table 2	
Comparison of 1995 and 2000 statistical models	9
Table 3	
Method of producing the 2000 estimates according to data source and type	9
Table 4	
Maternal mortality estimates by United Nations MDG regions, 2000	11
Table 5	
Comparison of 1995 and 2000 regional and global totals	12

Annex Tables

Annex Table A

Maternal mortality data derived from vital registration: Countries and territories with good death registration and good attribution of cause of death 16

Annex Table B

Maternal mortality data derived from vital registration: Countries with good death registration but uncertain attribution of cause of death 17

Annex Table C

Maternal mortality data derived from the direct sisterhood method: Reported and adjusted estimates 18

Annex Table D

Maternal mortality data derived from RAMOS 19

Annex Table E

Countries with data from household surveys or census 19

Annex Table F

Maternal mortality estimates derived from the model 20

Annex Table G

Estimates of number of maternal deaths, lifetime risk, maternal mortality ratio, and range of uncertainty (2000) 22

Annex Table H

Estimates of maternal mortality ratios, number of maternal deaths, and lifetime risk by WHO regions (2000) 27

Annex Table I

Estimates of maternal mortality ratios, number of maternal deaths, and lifetime risk by UNICEF regions (2000) 27

Annex Table J

Estimates of maternal mortality ratios, number of maternal deaths, and lifetime risk by UNFPA regions (2000) 28

Figures

Figure 1

Plot of residuals against fitted values 28

References 29



Executive summary

Reduction of maternal mortality is one of the major goals of several recent international conferences and has been included in the Millennium Development Goals (MDGs). However, because measuring maternal mortality is difficult and complex, reliable estimates of the dimensions of the problem are not generally available, and assessing progress towards the goal is difficult. In recent years, new ways of measuring maternal mortality have been developed, bearing in mind the needs and constraints of developing countries in particular. As a result, there is considerably more information available today than was the case even a few years ago. Nonetheless, problems of underreporting and misclassification are endemic to all methods, and estimates that are based on household surveys are subject to wide margins of uncertainty because of sample size issues. For all these reasons, it is difficult to compare the data obtained from different sources and to assess the overall magnitude of the problem.

In response to these challenges and in order to improve the information base, WHO, UNICEF and UNFPA have developed an approach to estimating maternal mortality that seeks both to generate estimates for countries with no data and to correct available data for underreporting and misclassification. A dual strategy is used which involves adjusting available country data and developing a simple model to generate estimates for countries without reliable information. The approach, with some variations, was used to develop estimates for maternal mortality in 1990 and 1995 and has been used again for generating these estimates for the year 2000.

On the basis of the present exercise, the estimated number of maternal deaths in 2000 for the world was 529,000 (Table 1). These deaths were almost equally divided between Africa (251,000) and Asia (253,000), with about 4% (22,000) occurring in Latin America and the Caribbean, and less than 1% (2,500) in the more developed regions of the world. In terms of the maternal mortality ratio (MMR), the world figure is estimated to be 400 per 100,000 live births. By region, the MMR was highest in Africa (830), followed by Asia (330), Oceania (240), Latin America and the Caribbean (190), and the developed countries (20).

The country with the highest estimated number of maternal deaths is India (136,000), followed by Nigeria (37,000), Pakistan (26,000), the Democratic Republic of the Congo and Ethiopia (24,000 each), the United Republic of Tanzania (21,000), Afghanistan (20,000), Bangladesh (16,000), Angola, China and Kenya (11,000 each), Indonesia and Uganda (10,000 each). These 13 countries account for 67% of all maternal deaths.

However, the number of maternal deaths is the product of the total number of births and obstetric risk per birth, described by the MMR. On a risk-per-birth basis, the list looks rather different. With the sole exception of Afghanistan, the countries with the highest MMRs are in Africa. The highest MMRs of 1,000 or greater, are, in order of magnitude, Sierra Leone (2,000), Afghanistan (1,900), Malawi (1,800), Angola (1,700), Niger (1,600), the United Republic of Tanzania (1,500), Rwanda (1,400), Mali (1,200), Central African Republic, Chad, Guinea-Bissau, Somalia and Zimbabwe (1,100 each).

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_30084



云报告
https://www.yunbaogao.cn

云报告
https://www.yunbaogao.cn

云报告
https://www.yunbaogao.cn