

Distribution: General  
WHO/CDD/SER/91.14  
Original: English

***DIVISION OF CHILD HEALTH AND DEVELOPMENT***

***INDICATORS FOR  
ASSESSING  
BREASTFEEDING  
PRACTICES***

Reprinted report of an Informal Meeting  
11 - 12 June 1991  
Geneva, Switzerland



**WORLD HEALTH ORGANIZATION  
GENEVA**

This document is not a formal publication of the World Health Organization (WHO), and all rights are reserved by the Organization. The document may, however, be freely reviewed, abstracted, reproduced and translated, in part or in whole, but not for sale nor for use in conjunction with commercial purposes.

The views expressed in documents by named authors are solely the responsibility of those authors.

**World Health Organization**

**1991**

## Indicators for assessing breastfeeding practices

An informal meeting convened by the WHO Division of Diarrhoeal and Acute Respiratory Disease Control on behalf of the Organization's Working Group on Infant Feeding<sup>1</sup> was held on 11-12 June 1991, at WHO headquarters in Geneva. The purpose of the meeting was to reach a consensus on the definitions of key breastfeeding indicators and specific methodologies for their measurement. In addition to the WHO participants, the meeting was attended by representatives of UNICEF, the United States Agency for International Development (USAID) and the Demographic Health Surveys (DHS) Program of the Institute for Resource Development/Macro International Inc., who had played an important role in developing the proposed indicators. The Swedish International Development Agency (SIDA) was also invited but was unable to send a participant. The participants are listed in Annex 1.

This report summarizes the discussion and consensus reached on breastfeeding indicators derived from household survey data. No consensus was reached on proposed breastfeeding indicators to be measured through enquiries at health facilities. It was agreed that this topic required further discussion, bearing in mind, for example, the monitoring of the "Ten steps to successful breastfeeding".<sup>2</sup>

The participating organizations and agencies committed themselves to promoting, adopting and implementing at global and country programme levels indicators measuring the following elements of feeding:

- **EXCLUSIVE BREASTFEEDING**
- **PREDOMINANT BREASTFEEDING**
- **TIMELY COMPLEMENTARY FEEDING**
- **CONTINUED BREASTFEEDING AT ONE AND TWO YEARS OF AGE**
- **BOTTLE FEEDING**

Precise definitions of these indicators are given in section 5 of the report. The preceding sections describe the rationale for their selection and for arriving at the definitions.

### 1. PURPOSE OF THE INDICATORS

The main purpose of developing breastfeeding indicators is to have a common set of measures to assess breastfeeding practices and evaluate the progress of promotional programmes. Indicators should be limited in number, relatively easy to measure and interpret, and operationally useful. The focus of the indicators should be on intra-country comparison, although the degree of comparability between countries is also of interest.

<sup>1</sup> The WHO Working Group on Infant Feeding in 1991 comprised representatives of the Food and Nutrition Programme, the Division of Diarrhoeal and Acute Respiratory Disease Control, the Maternal and Child Health Programme, the Office of the Legal Counsel and the Special Programme of Research, Development and Research Training in Human Reproduction.

<sup>2</sup> *Protecting, promoting and supporting breast-feeding: the special role of maternity services: A joint WHO/UNICEF statement.* World Health Organization, Geneva, 1989.

In addition to being suitable for use at the country level, breastfeeding indicators should be designed keeping in mind the monitoring requirements of the goals and policies outlined in the Innocenti Declaration<sup>3</sup> and reflected in the statements of the World Summit for Children.

Although the benefits of breastfeeding in terms of child survival are well known, changes in child mortality are difficult to measure and cannot easily be attributed to specific interventions. Attitudes towards breastfeeding, awareness of the importance of breastfeeding, and support to enable mothers to breastfeed are all important outcomes of promotional activities in health programmes, but they may also be difficult to measure and/or interpret and may not reflect actual practice. Indicators of current breastfeeding practices can be relatively easily measured and are sensitive to changes resulting from programme activities. It is clear that research studies and special evaluations may require the measurement of other parameters and the definition of other indicators than those listed in this document.

## 2. DEFINITIONS OF BREASTFEEDING CATEGORIES

The definitions of breastfeeding categories used in this report (all of which apply to the 24-hour period preceding the enquiry) are as follows:

- **EXCLUSIVE BREASTFEEDING:** the infant has received only breastmilk from his/her mother or a wet nurse, or expressed breastmilk, and no other liquids or solids with the exception of drops or syrups consisting of vitamins, mineral supplements or medicines.
- **PREDOMINANT BREASTFEEDING:** the infant's predominant source of nourishment has been breastmilk. However, the infant may also have received water and water-based drinks (sweetened and flavoured water, teas, infusions, etc.); fruit juice; Oral Rehydration Salts (ORS) solution; drop and syrup forms of vitamins, minerals and medicines; and ritual fluids (in limited quantities). With the exception of fruit juice and sugar-water, no food-based fluid is allowed under this definition.
- **EXCLUSIVE BREASTFEEDING AND PREDOMINANT BREASTFEEDING** together constitute **FULL BREASTFEEDING.**
- **BREASTFEEDING:** the child has received breastmilk (direct from the breast or expressed).
- **COMPLEMENTARY FEEDING:** the child has received both breastmilk and solid (or semi-solid) food.
- **BOTTLE-FEEDING:** the child has received liquid or semi-solid food from a bottle with a nipple/teat.

Although bottle-feeding is not strictly a "breastfeeding" category, it was considered essential to include it among the key indicators because of its impact on breastfeeding. The objective is to measure the prevalence of this *mode* of feeding, irrespective of the *content* of the feed - e.g., infants receiving breastmilk in a bottle are also included here.

The breastfeeding categories described above are essentially the same as those described in "Schema for the development of breastfeeding definitions" adopted by IGAB (Interagency Group for

---

<sup>3</sup> *Innocenti Declaration on the Protection, Promotion and Support of Breastfeeding*, adopted by participants in the WHO/UNICEF policymakers' meeting (co-sponsored by USAID and SIDA), Florence, Italy, 30 July to 1 August 1990.

Action on Breastfeeding) in 1988 and published in *Studies in Family Planning*, 21: 226-230, 1990, by M. Labbok and K. Krasovek. The differences are: (1) a change in terminology from "almost exclusive" to "predominant"; (2) the acceptance of drops or syrups in the category "exclusive breastfeeding"; and (3) the acceptance of certain liquids and ritual fluids, in limited amounts, in the category "predominant breastfeeding".

The criteria for the inclusion of infants in the above feeding categories used in developing the indicators are summarized in Table 1 below.

**Table 1: Criteria for inclusion in infant feeding categories**

<b>Category of infant-feeding</b>	<b>Requires that the infant receive</b>	<b>Allows the Infant to receive</b>	<b>Does not allow the infant to receive</b>
<b>Exclusive breastfeeding</b>	Breastmilk (including milk expressed or from wet nurse)	Drops, syrups (vitamins, minerals, medicines)	Anything else
<b>Predominant breastfeeding</b>	Breastmilk (including milk expressed or from wet nurse) as the predominant source of nourishment	Liquids (water, and water-based drinks, fruit juice, ORS), ritual fluids and drops or syrups (vitamins, minerals, medicines)	Anything else (in particular, non-human milk, food-based fluids)
<b>Complementary feeding</b>	Breastmilk and solid or semi-solid foods	Any food or liquid including non-human milk	
<b>Breastfeeding</b>	Breastmilk	"	
<b>Bottle-feeding</b>	Any liquid or semi-solid food from a bottle with nipple/teat	" Also allows breastmilk by bottle	

### **3. SELECTION OF AGE GROUPS FOR MEASURING BREASTFEEDING INDICATORS IN RELATION TO AGE-BASED FEEDING RECOMMENDATIONS**

Taking into consideration various policy documents, the meeting reached the following consensus on the interpretation of recommendations for infant and child feeding:

- "All infants should be fed exclusively on breastmilk from birth to 4-6 months of age": the inference of this statement is that 100% of infants up to exact age 4 months (< 120 days) should be exclusively breastfed.
- In order to meet their nutritional requirements, complementary foods should be introduced to the majority of infants during a transitional period lasting 2 months (that is, during the fifth and sixth months of life). Thus, nearly all infants older than exact age 6 months should be receiving complementary foods in addition to breastmilk.
- Children should be breastfed for at least one year and preferably for up to 2 years of age or beyond.

These recommendations are illustrated in the figure.

Taking into account the above feeding recommendations and the limitations of typical household surveys in terms of sample size, the meeting decided that, for the purpose of measuring indicators, four age groups should be used, and that, for the sake of simplicity, the four age groups should be of equal duration, i.e., 4 months each. These groups are also illustrated in the figure and defined below:

- **EXCLUSIVE BREASTFEEDING** and **PREDOMINANT BREASTFEEDING** are measured in infants up to exact age 4 months (< 120 days).
- **TIMELY COMPLEMENTARY FEEDING** is measured in infants older than exact age 6 months but less than exact age 10 months (180-299 days).
- **CONTINUED BREASTFEEDING** is measured twice, in children one year old, and in children nearing the end of their second year of life - that is, age groups older than exact age 12 months but less than exact age 16 months, and older than exact age 20 months but less than exact age 24 months.


#### 4. METHODOLOGY FOR MEASURING INDICATORS

The breastfeeding indicators derived from interviews at the household level will be measured using a household survey methodology. These indicators will be based on all live children less than 24 months of age (not yet having had their second birthday). Deceased children will not be included. The Innocenti Declaration recommends breastfeeding beyond 2 years; however, a sample of children less than 24 months will cover most of the period of interest and all of the above indicators. The indicators will be based on *current status data*, i.e., the current age of the child and other information for the 24 hours preceding the survey, rather than on retrospective data; mothers will not be asked when they stopped or started particular feeding practices, which are questions that tend to produce a heaping of data at certain ages. The 24-hour recall period for feeding practices was selected because it has been widely used and found appropriate in surveys of dietary intake.

Since it is the mother's behaviour *vis-à-vis* her child(ren) that is of interest, it is recommended that the estimates be based on all children born in a given time period, rather than including only the last-born child. If only last-born children are included in the sample, the findings may be biased, and the bias may not be equal in all countries or among all population subgroups. Last births are not a representative sample of all births. In addition, last births are spread out over an unspecified period. The sample of children for whom the indicator is calculated should always refer to all children born during a specified time period in order to be representative of a population of children.

A list of sample questions for use in surveys on breastfeeding indicators is presented in Annex 2.

**Figure: Age groups for measuring indicators in relation to feeding recommendations**



## 5. DEFINITIONS OF KEY INDICATORS

The key indicators are defined and explained below. A summary list of the indicators is presented in Annex 3.

### EXCLUSIVE BREASTFEEDING RATE

Proportion of infants less than 4 months of age who were exclusively breastfed:

$$= \frac{\text{Infants < 4 months (< 120 days) of age} \\ \text{who were exclusively breastfed in the last 24 hours}}{\text{Infants < 4 months (< 120 days) of age}}$$

#### Explanatory notes:

All infants should be fed exclusively on breastmilk from birth to 4-6 months of age. The exact age at which complementary feeding should be introduced will vary from child to child; however, implicit in the recommendation of the 4-6 months range is that all infants less than exact age 4 months (< 120 days) should be exclusively breastfed. Individual infants 120 days or older should be receiving complementary foods in addition to breastmilk if their growth on exclusive breastfeeding starts to falter.

This indicator includes breastfeeding from a wet nurse and feeding on expressed breastmilk. It was, however, thought simpler to retain the term "exclusive breastfeeding" rather than the more precise but cumbersome term "fed exclusively on breastmilk". (For the definition of "exclusive breastfeeding" see section 2 and Table 1.)

Using a 24-hour recall period may cause the proportion of exclusively breastfed infants to be slightly overestimated, since some infants who are given other liquids irregularly may not have received them in the 24 hours before the survey.

Although this rate may be low, at least initially, an increase in the proportion of exclusively breastfed infants will be the goal of many programmes, and thus it is desirable to have an indicator that measures the change.

### PREDOMINANT BREASTFEEDING RATE

Proportion of infants less than 4 months of age who are predominantly breastfed:

$$= \frac{\text{Infants < 4 months (< 120 days) of age} \\ \text{who were predominantly breastfed in the last 24 hours}}{\text{Infants < 4 months (< 120 days) of age}}$$

#### Explanatory notes:

As the proportion of infants exclusively breastfed may be very low, the intent of this indicator is to identify infants whose predominant source of nourishment is breastmilk, but who also receive other fluids. (For the definition of "predominant breastfeeding" see section 2 and Table 1.)